

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, England and Wales, 2001-2012, <sup>1,2,3,4,5,6,7</sup>

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	187.8	179.4	170.6	157.4	147.3	136.0	128.8	122.0	113.9	109.3	99.2	95.4
	2	Cerebrovascular diseases	69.2	69.3	66.5	59.9	55.7	52.5	48.7	47.4	43.9	43.0	35.7	34.1
	3	Malignant neoplasm of trachea, bronchus and lung	57.3	56.1	53.9	52.3	51.3	51.0	49.9	49.6	48.1	46.6	45.7	44.2
	4	Chronic lower respiratory diseases	42.9	42.4	43.6	39.4	39.2	36.6	36.1	37.1	34.5	33.7	34.0	35.1
	5	Influenza and pneumonia	38.9	39.0	40.9	36.4	35.7	32.4	30.8	30.8	29.3	27.0	27.3	26.1
	6	Malignant neoplasm of prostate	27.4	27.3	27.4	26.8	25.7	25.2	25.0	24.2	24.3	24.2	23.8	23.0
	7	Malignant neoplasm of colon, sigmoid, rectum and anus	24.4	24.2	23.6	23.5	23.0	22.3	21.7	22.0	21.1	21.2	20.3	20.5
	8	Dementia and Alzheimer's disease	15.4	15.4	15.4	14.6	14.3	14.3	15.9	17.0	16.6	17.6	27.9	31.3
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	18.5	18.8	18.8	17.8	17.3	17.6	17.4	17.0	16.7	16.1	16.3	16.5
	10	Aortic aneurysm and dissection	16.9	16.7	16.1	15.4	14.1	13.5	12.9	11.9	10.8	10.6	9.2	8.9
	18	Malignant neoplasm of bladder	9.3	9.0	8.8	8.5	8.1	8.1	8.1	8.0	8.0	7.6	7.7	7.6

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	88.0	84.6	81.4	73.8	68.8	63.2	59.5	56.5	50.8	48.2	43.4	42.6
	2	Cerebrovascular diseases	62.2	61.8	60.7	55.3	52.0	48.1	45.7	44.5	41.1	40.1	33.1	32.7
	3	Influenza and pneumonia	30.8	31.9	33.8	29.8	29.9	26.3	25.3	25.8	23.4	21.8	21.7	20.8
	4	Dementia and Alzheimer's disease	18.2	18.7	20.1	19.0	18.8	18.5	20.5	22.2	21.8	23.8	33.8	38.4
	5	Chronic lower respiratory diseases	24.9	25.4	27.4	24.2	25.1	24.0	24.4	25.3	23.2	23.9	24.1	25.2
	6	Malignant neoplasm of trachea, bronchus and lung	28.4	28.5	28.5	28.4	29.0	30.0	30.1	30.8	29.9	30.1	29.8	29.8
	7	Malignant neoplasms of breast	30.8	30.3	29.4	28.5	28.3	27.7	26.8	26.3	25.4	24.5	24.4	23.9
	8	Heart failure and complications and ill-defined heart disease	12.6	13.2	13.2	11.7	11.2	9.9	9.8	10.0	9.4	9.1	6.8	7.0
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	14.8	14.7	14.4	14.3	14.2	14.0	14.0	13.8	13.2	12.9	12.7	13.1
	10	Diseases of the urinary system	7.5	8.1	8.8	9.1	9.9	10.0	10.1	10.7	10.5	10.8	8.0	8.1
	29	Malignant neoplasm of bladder	2.9	3.0	3.0	2.8	2.8	2.9	2.7	2.7	2.7	2.6	2.6	2.7

1 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

2 Data for England and Wales includes deaths of non-residents

[3 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A in the following link](#)

4 Rankings are based on the grand total number of deaths for that cause of death

5 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

6 .. Rates were not calculated where there were less than 10 deaths in a year

7 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Greater London, 2001-2012,<sup>1,2,3,4,5,6,7,8</sup>

Sex	Rank	Cause	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	178.0	168.6	161.4	150.2	139.0	127.2	125.2	118.6	109.2	103.1	91.5	91.5
	2	Cerebrovascular diseases	65.8	63.2	61.3	56.7	51.1	49.8	44.6	44.1	41.6	40.7	34.3	32.9
	3	Malignant neoplasm of trachea, bronchus and lung	61.0	56.9	55.4	55.1	50.8	52.5	49.1	48.8	47.2	43.4	46.0	45.0
	4	Chronic lower respiratory diseases	46.2	43.8	46.2	41.6	42.9	38.4	37.3	38.1	35.4	32.6	34.7	36.7
	5	Influenza and pneumonia	50.9	47.1	50.3	42.2	40.1	34.5	32.9	34.5	30.3	27.5	27.1	28.9
	6	Malignant neoplasm of prostate	26.3	26.3	25.2	26.5	26.3	23.1	22.9	22.7	23.3	23.5	22.5	21.0
	7	Malignant neoplasm of colon, sigmoid, rectum and anus	22.6	23.1	22.1	20.2	19.2	20.5	19.5	19.7	20.3	18.3	17.5	19.0
	8	Malignant neoplasms of lymphoid, haematopoietic and related tissue	19.8	19.6	18.4	18.0	18.4	17.5	17.8	15.5	16.5	15.5	16.9	15.3
	9	Heart failure and complications and ill-defined heart disease	15.0	16.6	20.9	18.2	19.0	17.1	16.0	18.2	16.9	18.1	12.3	11.6
	10	Dementia and Alzheimer's disease	12.3	12.7	12.1	11.8	12.0	12.5	12.6	14.6	14.7	15.2	25.6	29.1
	19	Malignant neoplasm of bladder	8.8	8.7	8.0	7.6	8.2	7.9	7.4	7.6	7.8	6.3	7.3	7.1

Sex	Rank	Cause	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	80.5	78.4	76.4	69.3	64.7	56.8	56.4	55.9	49.7	45.2	40.5	40.1
	2	Cerebrovascular diseases	55.7	51.5	55.8	49.4	46.9	41.6	38.5	37.6	35.1	34.0	28.4	28.3
	3	Influenza and pneumonia	38.7	38.9	40.9	35.9	33.1	28.7	26.5	28.0	24.2	22.8	21.9	22.1
	4	Dementia and Alzheimer's disease	15.5	16.1	15.9	15.6	14.2	13.8	15.3	17.5	17.2	19.6	28.6	32.2
	5	Malignant neoplasm of trachea, bronchus and lung	28.2	28.0	28.2	27.5	29.1	27.5	30.2	28.1	28.3	27.7	27.4	27.5
	6	Chronic lower respiratory diseases	24.2	24.5	25.8	24.5	24.3	23.0	22.0	22.9	19.9	21.7	21.7	24.1
	7	Malignant neoplasms of breast	31.4	30.3	30.3	29.4	27.7	26.2	24.7	25.8	25.5	24.4	23.4	22.9
	8	Heart failure and complications and ill-defined heart disease	14.2	16.3	17.2	16.5	15.6	14.3	14.3	14.5	14.4	14.4	11.5	12.6
	9	Diseases of the urinary system	9.3	10.7	12.3	11.9	12.3	12.4	11.2	12.4	10.9	11.6	7.7	7.8
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	13.8	14.5	13.7	15.0	13.5	13.6	12.9	13.5	12.5	11.9	12.2	11.9
		31	Malignant neoplasm of bladder	2.5	2.4	2.9	2.7	2.2	2.6	2.4	2.4	2.2	2.1	2.4

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A in the following link

5 Rankings are based on the grand total number of deaths for that cause of death

6 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

7 .. Rates were not calculated where there were less than 10 deaths in a year

8 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Camden, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

Sex	Rank	Cause	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	197.2	188.3	173.3	179.5	180.6	140.5	136.8	133.6	114.0	100.0	84.0	74.0
	2	Malignant neoplasm of trachea, bronchus and lung	71.1	45.1	67.9	57.4	66.6	40.9	52.0	53.1	49.3	41.3	49.4	51.0
	3	Cerebrovascular diseases	66.1	54.3	55.5	64.4	66.1	39.9	45.1	32.9	39.9	30.5	32.8	28.8
	4	Chronic lower respiratory diseases	30.3	48.2	28.6	37.4	48.4	34.1	43.7	43.8	26.8	36.0	36.9	46.8
	5	Influenza and pneumonia	49.0	51.0	47.4	34.8	40.0	31.4	28.6	28.9	24.0	21.0	12.4	20.9
	6	Cirrhosis and other diseases of liver	37.4	42.3	32.4	26.7	20.9	37.7	32.3	32.8	..	19.5	21.3	12.6
	7	Malignant neoplasm of prostate	24.5	19.5	22.6	21.0	27.0	17.6	30.3	23.7	23.5	22.5	10.7	21.1
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	25.3	37.7	15.8	23.7	21.7	23.4	17.8	19.5	13.6	21.9	18.5	20.1
	9	Suicide and injury/poisoning of undetermined intent	11.3	13.5	13.7	18.4	11.7	12.6	15.1	9.7	11.2	..	8.1	..
	10	Dementia and Alzheimer's disease	..	12.5	..	21.5	..	9.7	..	16.3	..	11.9	26.7	18.7
	22*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	12.0	..	..	..	

Sex	Rank	Cause	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	75.3	77.7	91.8	63.6	71.6	63.1	64.1	51.4	34.4	38.2	31.7	25.5
	2	Cerebrovascular diseases	56.6	42.4	36.4	45.4	33.6	35.8	35.6	26.5	30.2	18.8	22.2	22.4
	3	Malignant neoplasm of trachea, bronchus and lung	27.3	31.5	22.9	40.8	33.4	34.4	33.7	24.9	28.7	27.3	32.2	15.7
	4	Influenza and pneumonia	32.1	33.5	21.3	18.0	19.0	14.6	17.5	20.3	18.6	12.3	15.8	13.1
	5	Chronic lower respiratory diseases	23.7	23.7	23.4	23.1	17.1	27.1	19.8	24.4	19.5	14.6	21.1	25.7
	6	Dementia and Alzheimer's disease	20.0	14.7	14.2	22.2	20.0	19.2	17.0	11.8	9.2	17.0	15.7	21.3
	7	Malignant neoplasms of breast	37.6	27.4	23.4	33.4	25.6	25.0	31.9	28.1	19.0	18.7	16.9	25.7
	8	Heart failure and complications and ill-defined heart disease	15.6	11.0	11.5	16.3	9.2	15.5	12.1	9.1	7.8	16.2	..	..
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	11.4	19.7	15.9	10.6	12.8	16.8	8.7	10.4	13.1	10.0	12.6	11.4
	10	Diseases of the urinary system	10.2	10.1	11.3	11.8	8.9	..	11.9	9.8	..	11.2	10.3	..
	30*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Hackney, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

		<i>Rate per 100,000 population</i>												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	194.8	180.6	212.7	171.8	161.6	181.6	165.4	150.3	131.4	131.5	105.0	103.3
	2	Malignant neoplasm of trachea, bronchus and lung	64.9	77.5	81.4	76.2	51.3	91.9	40.9	55.3	57.0	53.4	51.1	74.1
	3	Cerebrovascular diseases	55.7	60.3	62.3	66.1	46.8	60.1	50.1	47.5	46.1	52.4	48.2	41.5
	4	Chronic lower respiratory diseases	50.8	42.6	53.5	34.5	60.3	50.5	45.4	53.4	30.6	32.8	59.1	51.5
	5	Influenza and pneumonia	27.1	48.8	45.4	44.5	34.6	26.6	39.3	35.3	25.6	30.0	23.3	22.8
	6	Malignant neoplasm of prostate	29.1	24.6	31.0	36.0	30.4	24.6	17.2	17.0	18.7	23.7	21.2	22.2
	7	Cirrhosis and other diseases of liver	21.5	26.1	28.7	27.9	26.9	22.1	13.6	21.7	16.2	15.9	16.3	17.6
	8	Suicide and injury/poisoning of undetermined intent	17.3	8.3	9.2	10.9	8.9	11.6	9.2	19.6	12.7	13.7	6.2	6.2
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	23.6	30.6	..	17.1	19.1	23.3	15.2	18.7	14.3	20.6	20.3	27.5
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	14.7	22.2	19.4	19.0	26.5	18.1	..	13.5	21.4	15.7	..	19.3
	22	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	14.5

		<i>Rate per 100,000 population</i>												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	103.3	99.9	80.1	75.7	71.1	69.1	65.2	81.8	72.9	65.5	58.6	54.7
	2	Cerebrovascular diseases	55.8	52.9	46.6	53.5	45.6	57.0	27.4	40.7	41.5	38.4	26.2	23.3
	3	Influenza and pneumonia	32.1	22.3	31.3	31.5	35.4	23.6	20.5	28.2	24.1	24.8	21.6	26.6
	4	Malignant neoplasm of trachea, bronchus and lung	29.0	27.2	25.8	27.5	27.8	30.4	35.5	34.6	40.6	37.0	32.3	32.9
	5	Chronic lower respiratory diseases	26.2	27.9	29.5	24.5	29.5	28.2	34.5	25.7	19.3	34.6	30.0	27.2
	6	Malignant neoplasms of breast	26.9	33.8	31.0	33.6	22.4	27.2	23.1	29.9	17.4	16.3	19.2	14.0
	7	Dementia and Alzheimer's disease	8.0	12.2	16.2	13.3	10.9	8.3	17.7	13.4	13.9	13.6	32.0	24.3
	8	Diseases of the urinary system	..	18.8	10.3	16.9	9.2	18.5	13.3	12.2	16.1	12.3	9.0	..
	9	Heart failure and complications and ill-defined heart disease	15.8	12.2	10.2	13.9	9.7	13.7	12.1	9.6	9.8	11.2	9.9	..
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	..	..	14.5	..	..	13.1	15.1	15.6	..	15.0	13.5	13.2
	28	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Hammersmith and Fulham, 2001-2012,<sup>1,2,3,4,5,6,7,8,9,10,11</sup>

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	153.8	134.3	154.4	140.6	132.5	98.4	118.4	114.5	83.7	98.8	113.9	90.6
	2	Malignant neoplasm of trachea, bronchus and lung	73.8	72.2	80.0	56.9	52.3	50.5	53.7	47.4	49.4	48.6	49.2	50.8
	3	Chronic lower respiratory diseases	66.2	34.3	60.9	47.5	54.7	32.1	55.7	42.3	57.5	34.4	48.9	44.4
	4	Cerebrovascular diseases	63.8	56.6	73.8	46.5	35.7	41.2	34.8	32.4	28.2	25.0	43.3	43.5
	5	Influenza and pneumonia	41.4	50.9	59.4	40.5	39.1	30.7	30.6	36.0	22.3	24.9	26.1	29.5
	6	Cirrhosis and other diseases of liver	26.5	25.8	27.9	35.4	30.6	30.0	21.2	22.6	27.5	18.4	28.1	19.3
	7	Malignant neoplasm of prostate	28.6	18.3	24.4	34.5	19.0	14.0	28.6	..	28.7	18.0	31.5	..
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	..	25.5	25.2	18.8	22.4	19.8	22.4	23.0	22.6	19.3	26.5	23.8
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	25.7	19.3	20.9	15.6	28.7	16.9	27.5	16.5	18.3	..	14.7	15.9
	10	Suicide and injury/poisoning of undetermined intent	14.5	10.2	..	14.3	16	9.4	7.2	16	9.8	9.7	8.6	14.5
19	Malignant neoplasm of bladder	..	..	..	..	..	16.1	..	15.2	..	..	..	..	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	71.0	68.4	54.1	63.2	56.4	59.7	48.9	53.1	42.7	44.1	43.3	37.3
	2	Cerebrovascular diseases	55.9	40.5	39.0	39.8	37.4	43.4	39.7	40.8	33.8	27.9	31.4	20.4
	3	Influenza and pneumonia	34.9	30.4	35.8	29.9	27.0	20.6	23.8	20.5	18.6	22.5	22.6	15.8
	4	Malignant neoplasm of trachea, bronchus and lung	28.3	46.5	35.2	36.8	46.2	26.9	27.2	43.6	26.3	31.5	43.7	36.5
	5	Chronic lower respiratory diseases	23.5	19.2	20.7	25.9	29.6	34.4	16.6	33.3	24.1	34.1	25.5	26.2
	6	Malignant neoplasms of breast	31.1	27.4	37.2	26.7	24.3	17.8	23.8	24.4	25.0	23.3	27.5	22.6
	7	Dementia and Alzheimer's disease	7.4	..	7.4	..	7.5	10.7	11.2	15.5	16.9	26.5	38.1	37.0
	8	Heart failure and complications and ill-defined heart disease	8.7	14.1	22.9	15.3	19.2	8.2	11.1	13.4	9.6	7.9	..	10.2
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	14.9	21.9	10.5	12.4	16.1	14.9	5.6	17.7	17.3	11.2	18.1	15.8
	10*	Diseases of the urinary system	..	7.5	..	9.7	..	..	7.5	15.5	..	9.8	..	..
	10*	Malignant neoplasms of lymphoid, haematopoietic and related tissue	11.5	..	12.4	13.2	..	10.9	14.1	9.8	..	..	..	..
29*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Haringey, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	189.1	209.8	200.8	162.1	157.7	119.7	160.1	144.3	113.6	108.5	87.4	90.6
	2	Cerebrovascular diseases	76.0	96.9	76.2	74.0	57.1	63.6	51.8	47.1	45.9	41.5	26.9	39.7
	3	Malignant neoplasm of trachea, bronchus and lung	63.3	56.2	47.5	60.7	48.1	70.8	46.2	47.0	54.8	32.0	51.6	56.3
	4	Influenza and pneumonia	59.1	55.7	69.6	55.9	33.1	30.3	37.6	30.9	32.6	18.3	20.7	30.0
	5	Chronic lower respiratory diseases	58.8	55.7	39.5	44.9	41.3	34.7	39.5	25.6	23.8	28.4	20.4	36.9
	6	Malignant neoplasm of prostate	33.2	28.6	32.8	26.8	25.8	13.7	40.3	21.4	31.9	26.6	28.4	18.8
	7	Cirrhosis and other diseases of liver	16.9	20.5	16.9	24.6	27.2	27.6	25.4	17.9	23.6	28.4	14.2	16.8
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	21.6	13.9	15.8	23.6	15.2	27.5	22.6	25.0	17.0	20.6	21.6	19.6
	9	Suicide and injury/poisoning of undetermined intent	14.8	12.9	12.5	10	15.1	9.3	9.6	13.5	11.8	12.2	8.3	13.1
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	18.1	22.6	20.8	20.9	18.0	21.4	19.9	20.3	14.9	21.0	16.4	..
	21	Malignant neoplasm of bladder	..	..	14.5	..	..	..	14.9	..	..	..	12.6	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	79.4	79.6	85.5	67.5	60.2	64.7	58.9	59.8	57.7	59.7	40.1	36.8
	2	Cerebrovascular diseases	64.7	58.0	59.4	52.3	42.1	42.2	47.5	26.2	30.6	32.4	28.1	31.0
	3	Influenza and pneumonia	46.4	50.8	42.6	37.9	34.7	24.1	22.5	25.8	16.1	23.1	19.6	23.4
	4	Malignant neoplasms of breast	36.4	22.2	38.5	28.2	35.8	20.9	29.2	30.8	29.5	27.2	27.0	29.6
	5	Malignant neoplasm of trachea, bronchus and lung	27.4	26.4	23.1	23.7	28.4	16.8	26.9	25.4	22.0	35.3	28.9	18.2
	6	Heart failure and complications and ill-defined heart disease	21.2	19.7	22.9	20.1	10.2	16.6	15.2	13.6	19.8	12.4	20.2	25.5
	7	Chronic lower respiratory diseases	17.6	20.2	21.0	20.3	21.9	15.6	23.7	15.2	17.8	13.4	14.7	22.8
	8	Dementia and Alzheimer's disease	12.7	18.0	14.3	10.8	12.1	14.0	13.9	..	9.4	17.0	21.9	18.9
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	25.2	11.5	16.7	16.3	..	14.4	17.5	12.3	14.6	20.9	12.2	16.1
	10	Diseases of the urinary system	14.7	16.6	14.7	13.6	7.6	7.5	8.6	8.3	..	10.5	..	7.1
	35*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Islington, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

Sex	Rank	Group description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	253.9	202.7	176.6	226.8	165.7	164.7	171.4	141.6	157.5	117.8	121.9	94.4
	2	Malignant neoplasm of trachea, bronchus and lung	99.1	92.5	83.2	76.5	87.5	73.7	59.8	77.6	58.8	56.0	64.6	62.8
	3	Cerebrovascular diseases	63.9	53.3	79.5	41.9	46.6	68.3	62.5	42.7	54.7	47.3	41.3	33.4
	4	Chronic lower respiratory diseases	75.2	50.4	63.1	48.0	53.7	35.4	29.4	57.8	54.0	37.5	46.6	49.6
	5	Influenza and pneumonia	57.2	54.8	44.1	42.7	41.8	39.7	43.8	51.0	25.9	26.5	35.1	47.3
	6	Cirrhosis and other diseases of liver	23.0	31.7	22.1	..	22.8	34.1	23.9	24.0	23.9	31.9	22.1	17.5
	7	Malignant neoplasm of prostate	29.5	32.9	26.1	26.4	23.7	25.8	15.0	21.1	23.9	22.7	20.5	22.8
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	32.0	32.4	26.7	22.8	..	23.7	19.9	21.2	32.8	19.5	17.8	30.5
	9	Suicide and injury/poisoning of undetermined intent	13	18.5	12.4	22.3	19.2	13.5	9.8	9.7	11.7	15.9	7.6	..
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	18.6	..	..	33.1	20.3	16.7	23.5	..	16.5	11.8	24.5	15.6
	19	Malignant neoplasm of bladder	..	18.2	..	16.0	..	..	..	..	..	..	..	

Sex	Rank	Group description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	96.4	81.6	108.5	86.9	85.9	68.7	72.6	60.3	67.3	43.8	28.5	51.0
	2	Cerebrovascular diseases	58.7	48.4	51.1	41.9	64.7	36.5	57.8	43.1	30.4	27.1	29.6	26.8
	3	Malignant neoplasm of trachea, bronchus and lung	34.7	48.0	31.8	43.7	39.7	40.6	42.1	34.3	42.5	42.4	57.6	33.5
	4	Influenza and pneumonia	38.5	31.9	33.0	37.7	42.9	26.6	22.1	23.8	25.3	21.1	21.8	18.3
	5	Chronic lower respiratory diseases	27.0	40.4	25.3	37.5	28.5	25.7	35.5	36.7	22.6	22.6	28.4	25.7
	6	Malignant neoplasms of breast	29.3	32.0	31.9	31.0	20.2	31.4	29.6	26.0	36.1	33.6	19.1	12.8
	7	Dementia and Alzheimer's disease	19.6	12.5	14.9	18.2	14.8	16.8	15.9	17.9	14.5	18.2	27.7	35.8
	8	Heart failure and complications and ill-defined heart disease	13.9	20.7	13.4	20.0	15.9	17.2	9.3	12.7	..	11.6	..	..
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	19.5	19.1	11.3	12.9	16.6	13.1	12.6	13.7	20.3	8.7	12.7	9.1
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	11.7	19.0	18.6	15.2	..	10.4	12.5	..	11.3	11.7	11.8	14.4
	32*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Kensington and Chelsea, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

		Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	124.4	95.3	120.4	98.4	96.0	84.0	75.7	77.6	68.1	64.6	49.3	48.4
	2	Malignant neoplasm of trachea, bronchus and lung	41.5	65.3	31.4	25.8	38.8	44.7	40.2	35.8	47.6	29.3	48.6	41.4
	3	Cerebrovascular diseases	34.9	50.0	47.7	38.7	27.1	31.4	27.3	37.4	39.0	43.2	37.1	27.4
	4	Chronic lower respiratory diseases	34.4	31.0	34.4	40.1	32.8	28.0	28.2	27.0	18.6	27.7	35.0	41.0
	5	Influenza and pneumonia	31.5	22.1	17.9	21.5	22.3	23.6	23.8	16.0	12.4	23.7	25.2	23.1
	6	Malignant neoplasm of prostate	20.4	18.7	21.5	16.9	14.7	18.8	16.8	16.2	20.3	22.9	15.3	17.7
	7	Heart failure and complications and ill-defined heart disease	13.6	15.9	16.8	22.4	21.5	14.3	19.4	28.9	13.0	19.3	18.2	10.0
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	25.0	17.5	20.0	19.9	17.1	..	17.9	18.8	19.2	21.2	13.0	15.2
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	18.2	13.4	19.7	16.0	20.4	13.2	19.4	13.4	13.0	15.4	11.2	11.6
	10	Dementia and Alzheimer's disease	..	..	..	11.4	10.3	..	11.9	11.5	10.4	..	25.0	25.0
	17	Malignant neoplasm of bladder	..	..	..	..	..	..	..	15.7	..	..	..	

		Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	48.6	54.6	40.7	42.8	43.1	30.6	47.7	35.3	28.5	30.9	31.9	30.5
	2	Cerebrovascular diseases	33.3	25.6	35.3	39.4	32.1	23.5	26.0	26.7	27.7	20.4	27.1	24.1
	3	Influenza and pneumonia	22.0	27.6	18.7	23.6	22.4	25.1	15.1	17.3	10.7	15.2	12.8	18.2
	4	Malignant neoplasm of trachea, bronchus and lung	19.7	16.6	29.7	22.1	28.1	14.2	22.6	24.6	25.3	24.2	31.4	26.9
	5	Malignant neoplasms of breast	23.4	24.7	15.9	22.1	16.0	22.1	22.1	20.1	24.6	24.2	30.6	21.7
	6	Chronic lower respiratory diseases	19.5	21.5	21.3	20.4	18.0	20.0	11.8	18.4	20.4	17.5	14.3	19.9
	7	Dementia and Alzheimer's disease	10.4	13.2	13.4	8.2	11.7	..	9.2	8.4	10.9	15.2	21.0	23.2
	8	Heart failure and complications and ill-defined heart disease	9.8	9.5	6.8	17.6	15.3	10.3	15.9	14.2	10.0	13.0	10.7	11.5
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	13.5	10.8	11.1	9.6	12.8	..	14.8	15.0	13.2	..	14.6	13.0
	10	Diseases of the urinary system	..	9.0	..	..	6.2	8.8	8.2	7.6	8.0	..	..	10.6
	37*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics



Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Lambeth, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

		<i>Rate per 100,000 population</i>												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	194.0	173.0	147.4	164.9	139.7	152.9	141.6	129.0	115.7	105.4	80.3	105.4
	2	Malignant neoplasm of trachea, bronchus and lung	87.5	83.5	62.1	63.6	71.1	68.0	62.3	74.2	69.8	54.9	64.7	54.7
	3	Cerebrovascular diseases	87.6	65.1	73.2	74.8	81.9	67.0	46.5	50.7	41.1	43.5	36.4	40.0
	4	Chronic lower respiratory diseases	56.5	62.4	61.0	64.1	54.0	65.4	46.3	46.8	41.4	41.0	51.6	46.9
	5	Influenza and pneumonia	43.8	55.4	52.7	31.9	34.4	33.8	32.2	30.2	31.8	16.9	28.0	32.6
	6	Malignant neoplasm of prostate	23.2	37.8	21.7	21.0	27.8	39.0	24.5	34.9	34.4	25.8	25.0	27.7
	7	Cirrhosis and other diseases of liver	29.5	26.2	26.8	27.5	27.5	31.2	35.1	21.5	15.4	14.3	17.1	14.2
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	19.6	26.2	25.2	22.7	20.6	21.9	20.8	18.2	19.5	20.1	14.0	14.3
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	26.5	22.5	14.1	14.7	19.5	11.0	14.8	20.0	22.8	23.2	20.9	15.5
	10	Suicide and injury/poisoning of undetermined intent	12.5	7.6	13.9	11.7	9.4	12.9	8.9	9.3	11.7	..	10.4	8.7
	26	Malignant neoplasm of bladder	11.0	..	..	..	..	..	..	..	..	..	..	..

		<i>Rate per 100,000 population</i>												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	83.8	83.1	76.3	86.5	64.2	60.7	53.3	57.1	60.4	50.3	38.3	36.5
	2	Cerebrovascular diseases	59.3	50.6	54.3	71.1	69.2	53.6	39.4	46.3	43.1	34.7	17.9	29.8
	3	Influenza and pneumonia	31.1	30.0	38.2	29.4	27.2	21.5	20.7	22.0	25.3	23.7	29.8	15.3
	4	Malignant neoplasm of trachea, bronchus and lung	34.7	41.6	34.6	28.0	46.4	34.4	36.8	26.4	39.5	28.2	26.0	37.0
	5	Chronic lower respiratory diseases	35.2	28.6	33.7	27.4	29.4	18.2	33.2	31.0	29.7	28.4	31.8	29.6
	6	Dementia and Alzheimer's disease	24.0	20.0	14.4	18.0	20.4	16.7	18.8	20.8	22.1	28.9	27.4	34.5
	7	Malignant neoplasms of breast	31.0	33.8	28.2	28.8	30.5	21.8	25.1	30.1	24.4	26.7	22.6	21.3
	8	Diseases of the urinary system	7.7	11.5	12.4	16.4	16.0	12.9	11.8	15.2	20.5	14.7	10.1	9.7
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	10.5	9.7	18.1	15.5	14.7	8.5	17.4	16.8	9.7	12.3	14.1	17.9
	10	Heart failure and complications and ill-defined heart disease	15.1	9.3	16.6	12.0	12.1	10.1	9.9	6.9	13.3	..	..	..
	34	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	..

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Lewisham, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	190.4	175.3	170.8	179.6	173.5	166.6	163.0	146.4	143.4	118.6	85.1	92.7
	2	Cerebrovascular diseases	80.1	70.4	66.8	78.2	69.6	39.3	59.6	55.1	56.2	46.4	43.8	35.4
	3	Malignant neoplasm of trachea, bronchus and lung	66.1	60.7	67.1	62.0	67.8	62.4	60.8	59.0	55.7	52.4	70.3	49.7
	4	Chronic lower respiratory diseases	77.2	68.1	62.4	65.7	63.2	46.1	57.9	54.6	46.1	44.3	38.3	52.1
	5	Influenza and pneumonia	80.2	50.8	68.8	61.4	44.5	41.0	37.1	61.7	30.2	32.8	43.0	30.8
	6	Malignant neoplasm of prostate	35.8	31.0	35.7	32.3	30.2	23.5	24.1	26.2	22.5	21.2	27.1	22.0
	7	Malignant neoplasm of colon, sigmoid, rectum and anus	35.0	19.9	27.9	23.9	16.2	21.8	21.9	21.6	29.3	19.4	28.8	18.3
	8	Dementia and Alzheimer's disease	9.2	22.3	18.0	16.4	21.0	10.4	11.2	19.7	14.8	21.7	30.9	46.8
	9	Cirrhosis and other diseases of liver	28.2	26.2	23.9	24.5	23.8	18.4	14.4	23.1	16.9	20.5	14.6	21.9
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	27.0	18.1	18.7	18.9	13.0	14.2	12.5	21.7	15.2	17.2	17.8	22.8
	20	Malignant neoplasm of bladder	..	..	12.5	11.7	..	..	..	..	9.8	..	9.5	..

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	91.5	93.0	99.6	83.1	99.6	78.5	72.9	75.9	58.7	48.4	47.6	42.8
	2	Cerebrovascular diseases	60.6	64.6	60.5	61.9	52.6	52.7	49.0	47.6	45.8	50.8	28.4	34.6
	3	Influenza and pneumonia	53.5	56.0	61.1	58.0	41.8	38.1	33.9	36.2	27.8	29.3	26.8	23.2
	4	Dementia and Alzheimer's disease	18.0	24.4	24.0	18.6	22.6	15.8	23.2	23.2	18.9	19.2	46.5	31.7
	5	Chronic lower respiratory diseases	33.7	25.8	33.5	30.9	33.5	26.3	26.7	38.2	21.3	28.2	27.8	21.5
	6	Malignant neoplasm of trachea, bronchus and lung	38.9	35.1	31.2	36.2	28.7	38.8	32.9	35.9	28.6	27.2	34.8	25.6
	7	Malignant neoplasms of breast	31.6	36.8	36.7	31.7	34.8	26.0	23.8	25.4	25.6	20.5	26.0	27.4
	8	Diseases of the urinary system	14.7	11.4	16.7	9.2	19.0	20.9	17.0	17.7	13.5	17.0	14.3	12.6
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	12.4	12.0	17.0	19.0	13.6	14.9	12.7	14.6	14.4	13.3	14.8	12.9
	10	Heart failure and complications and ill-defined heart disease	7.6	14.1	6.7	6.8	..	5.9	7.2	8.0	5.8	17.4	14.7	15.6
	30*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Newham, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	258.3	222.7	178.8	172.4	190.4	158.7	133.2	169.2	125.6	147.2	102.7	124.0
	2	Cerebrovascular diseases	71.2	83.8	70.6	56.0	59.7	58.3	64.8	75.8	60.7	47.2	39.2	45.0
	3	Malignant neoplasm of trachea, bronchus and lung	83.3	73.5	60.4	72.5	72.7	64.6	46.2	55.2	57.8	52.8	59.1	55.0
	4	Chronic lower respiratory diseases	62.8	62.8	44.6	44.2	55.9	64.4	46.6	56.4	58.1	49.1	39.1	61.9
	5	Influenza and pneumonia	71.5	48.0	60.4	54.1	61.9	41.5	43.5	34.0	37.5	31.0	26.4	28.5
	6	Cirrhosis and other diseases of liver	24.4	31.1	29.3	30.0	20.9	24.6	19.5	23.8	24.5	23.1	25.3	14.0
	7	Malignant neoplasm of prostate	24.5	26.1	21.5	29.5	20.2	25.8	28.5	36.0	22.5	25.1	18.8	28.9
	8	Dementia and Alzheimer's disease	..	13.7	18.3	27.6	14.3	18.4	22.8	20.1	14.9	15.9	28.8	40.9
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	24.5	26.4	20.0	17.7	19.8	26.0	16.1	25.9	23.4	14.0	13.5	21.2
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	35.0	..	14.4	20.5	..	20.4	12.3	16.9	20.3	12.0	17.6	10.3
18	Malignant neoplasm of bladder	16.6	11.1	..	..	..	15.0	..	..	..	..	..	..	

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	93.0	108.4	98.3	93.2	103.6	70.6	93.9	64.6	57.9	67.8	46.5	44.5
	2	Cerebrovascular diseases	68.4	65.5	61.5	56.3	73.2	56.4	49.9	42.6	48.1	47.6	32.0	28.6
	3	Influenza and pneumonia	49.3	58.2	70.1	48.2	46.0	34.9	31.8	28.1	25.6	24.1	26.0	28.5
	4	Malignant neoplasm of trachea, bronchus and lung	38.9	30.2	32.4	41.4	37.4	42.9	44.5	25.9	33.6	29.1	27.6	34.2
	5	Chronic lower respiratory diseases	23.2	23.8	31.5	37.3	35.4	37.3	24.7	30.2	30.5	31.3	30.5	32.9
	6	Dementia and Alzheimer's disease	17.5	22.4	25.3	19.9	19.2	18.5	19.1	21.9	23.3	20.5	32.0	40.3
	7	Malignant neoplasms of breast	41.1	20.0	31.3	36.9	17.9	32.1	21.2	22.9	28.1	23.6	23.4	26.5
	8	Diseases of the urinary system	14.9	12.3	20.5	13.0	18.5	18.0	23.1	15.5	15.5	22.0	12.1	8.0
	9	Heart failure and complications and ill-defined heart disease	12.4	15.8	11.5	14.1	23.1	17.3	17.6	13.9	17.1	12.1	..	10.8
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	20.2	14.6	12.7	10.1	11.0	15.5	11.5	13.2	13.2	..	15.2	13.1
29	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

[4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Southwark, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

		Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	180.3	173.7	147.3	143.9	158.6	120.1	130.3	129.7	102.9	102.1	114.2	89.3
	2	Malignant neoplasm of trachea, bronchus and lung	67.8	78.2	81.4	77.9	65.9	60.8	61.4	80.5	70.7	61.9	54.1	56.9
	3	Chronic lower respiratory diseases	53.2	59.6	68.7	50.5	79.8	63.9	59.9	65.6	55.5	55.9	65.6	47.4
	4	Cerebrovascular diseases	69.5	71.4	64.9	55.9	43.6	54.4	50.3	48.6	39.7	43.3	36.3	34.8
	5	Influenza and pneumonia	46.1	46.3	52.0	41.5	33.7	31.3	29.6	38.1	31.1	23.3	35.7	28.0
	6	Cirrhosis and other diseases of liver	28.1	22.5	28.3	19.5	25.9	32.4	32.6	20.8	20.9	16.3	20.9	15.8
	7	Malignant neoplasm of prostate	15.7	22.4	29.6	12.4	18.9	9.9	20.5	20.0	32.3	26.0	21.0	22.2
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	14.4	20.3	19.4	18.9	37.5	22.9	18.8	26.5	21.1	24.1	12.8	21.3
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	21.3	26.1	24.4	15.9	18.6	24.9	22.4	20.5	13.5	18.8	19.9	14.2
	10	Suicide and injury/poisoning of undetermined intent	8.4	11.2	14.2	11.5	11.7	9.3	10.7	9	7	10.2	11	10.4
	21	Malignant neoplasm of bladder	..	..	..	..	10.2	..	..	..	..	..	..	13.3

		Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	85.2	76.7	82.1	76.6	60.6	63.8	65.2	48.0	42.8	54.5	38.1	39.2
	2	Cerebrovascular diseases	47.2	53.0	48.6	51.2	51.6	30.3	40.5	34.7	40.2	34.3	24.4	26.5
	3	Chronic lower respiratory diseases	31.4	43.9	32.5	36.2	37.9	36.3	39.3	31.0	29.8	37.1	35.6	26.2
	4	Influenza and pneumonia	37.5	35.2	27.5	24.5	34.4	21.5	22.3	26.5	21.0	22.9	23.7	17.7
	5	Malignant neoplasm of trachea, bronchus and lung	35.7	33.0	34.5	38.1	40.4	30.1	39.7	46.7	30.4	31.8	39.4	30.6
	6	Dementia and Alzheimer's disease	9.5	16.2	14.8	17.1	12.5	13.1	14.3	15.8	20.0	21.9	30.4	38.4
	7	Malignant neoplasms of breast	19.3	32.6	25.6	26.9	26.0	31.1	22.5	31.4	23.6	24.8	24.5	21.1
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	18.5	14.8	9.9	15.4	15.4	20.6	13.8	19.0	12.1	20.6	12.9	15.7
	9	Diseases of the urinary system	9.2	8.1	16.3	10.6	12.5	13.9	11.9	14.4	14.7	11.5	7.1	..
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	10.8	16.3	11.6	7.7	14.9	9.3	10.6	14.3	6.9	9.3	..	17.7
	25*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Tower Hamlets, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	221.0	245.9	231.0	188.0	189.0	193.8	152.3	193.8	157.1	137.3	113.4	130.5
	2	Malignant neoplasm of trachea, bronchus and lung	81.6	117.4	90.8	85.1	69.7	85.7	74.4	73.3	76.2	72.6	73.4	58.0
	3	Chronic lower respiratory diseases	76.7	63.1	62.2	54.9	81.9	77.6	54.7	69.3	55.7	53.0	57.6	63.3
	4	Cerebrovascular diseases	68.6	70.3	64.0	74.5	64.9	55.7	58.3	55.2	55.4	46.2	42.7	56.3
	5	Influenza and pneumonia	59.3	40.4	47.8	26.8	39.5	40.2	23.7	36.9	..	24.5	32.4	27.6
	6	Cirrhosis and other diseases of liver	23.4	28.0	30.0	25.0	20.7	26.0	16.2	17.9	17.1	22.1	26.7	21.5
	7	Suicide and injury/poisoning of undetermined intent	13.9	11.9	12.8	9.4	..	12	19.2	9.3	7.5	10.4	5.8	10.8
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	32.2	26.9	23.2	16.7	17.6	27.5	16.2	27.1	24.7	16.8	..	21.1
	9	Malignant neoplasm of prostate	28.0	28.9	16.3	26.3	16.9	19.4	18.8	16.4	21.3	13.8	19.9	19.8
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	20.2	19.6	14.0	..	21.5	..	15.9	16.3	15.9	..	25.0	..
21	Malignant neoplasm of bladder	13.7	..	..	..	..	..	..	..	..	13.9	..	..	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	112.4	108.9	97.1	106.0	81.9	81.8	75.4	64.8	57.4	62.6	58.7	54.8
	2	Cerebrovascular diseases	47.0	60.3	65.3	51.1	51.1	50.8	49.3	45.6	29.0	32.2	31.0	36.9
	3	Chronic lower respiratory diseases	42.2	35.0	52.8	39.7	47.9	44.9	32.4	44.4	32.9	49.2	45.5	31.6
	4	Malignant neoplasm of trachea, bronchus and lung	48.7	45.3	36.6	42.3	44.7	62.1	38.1	47.9	44.4	32.3	42.2	47.7
	5	Influenza and pneumonia	37.6	38.6	35.5	24.7	31.0	30.8	28.9	28.7	20.4	21.0	18.0	18.7
	6	Dementia and Alzheimer's disease	18.4	11.0	12.3	20.2	17.3	16.4	19.2	19.0	16.5	19.1	26.3	33.1
	7	Malignant neoplasms of breast	33.5	22.9	29.3	36.7	25	15.9	22.9	27.4	30.9	22.4	15.1	16.3
	8	Diseases of the urinary system	11.0	14.4	16.6	16.6	19.6	12.3	17.3	18.8	13.3	13.4	..	10.4
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	12.2	17.9	17.1	18.4	12.7	12.0	20.9	12.6	15.9	11.5	13.6	25.4
	10	Heart failure and complications and ill-defined heart disease	14.3	10.1	11.9	15.9	12.7	14.5	..	10.9	17.7	11.8	..	..
31*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Wandsworth, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

			Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	
Males	1	Ischaemic heart diseases	184.2	159.3	142.8	131.8	139.9	101.6	121.9	111.2	118.7	80.6	83.3	103.0	
	2	Malignant neoplasm of trachea, bronchus and lung	65.8	66.4	57.7	63.7	60.1	59.1	54.7	66.8	58.8	34.2	51.2	40.7	
	3	Cerebrovascular diseases	75.8	63.3	57.9	57.8	50.0	53.5	45.4	51.9	45.4	58.9	31.8	27.5	
	4	Chronic lower respiratory diseases	46.2	51.7	46.7	52.1	45.5	38.1	40.4	41.0	37.7	38.1	44.1	36.1	
	5	Influenza and pneumonia	58.8	39.6	65.2	42.6	47.1	37.9	41.4	25.9	32.7	22.9	20.4	35.1	
	6	Heart failure and complications and ill-defined heart disease	15.1	17.0	33.1	40.5	50.8	47.1	38.5	33.4	32.5	40.2	29.1	15.4	
	7	Malignant neoplasm of prostate	26.2	27.4	23.6	30.7	28.8	27.3	22.0	20.1	27.7	28.3	26.0	22.4	
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	23.6	27.3	16.5	22.2	29.0	17.3	18.0	26.5	21.3	22.2	19.8	15.6	
	9	Dementia and Alzheimer's disease	21.2	9.5	14.2	10.7	11.4	17.6	17.0	15.2	14.9	16.9	24.4	31.5	
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	23.9	20.8	16.9	21.3	20.3	17.6	16.8	15.4	16.1	18.3	11.6	23.5	
20	Malignant neoplasm of bladder	9.2	9.3	12.5	..	11.2	..	..	..	..	..	11.3	..		

			Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	
Females	1	Ischaemic heart diseases	83.8	88.3	79.5	70.0	68.2	56.8	55.0	63.6	54.3	32.8	42.1	50.0	
	2	Cerebrovascular diseases	54.7	63.5	57.2	49.6	53.4	39.6	37.3	43.8	32.4	37.3	22.3	27.8	
	3	Influenza and pneumonia	44.0	40.0	48.4	39.6	35.5	43.1	32.8	25.6	22.7	20.9	27.4	23.6	
	4	Dementia and Alzheimer's disease	25.7	19.9	14.2	17.3	18.4	11.7	18.8	19.5	21.7	18.9	36.7	30.8	
	5	Chronic lower respiratory diseases	38.2	29.1	33.0	31.3	26.7	26.2	26.7	28.6	22.6	27.5	22.9	22.9	
	6	Heart failure and complications and ill-defined heart disease	16.8	17.5	23.6	23.0	27.4	24.6	25.6	26.5	19.6	16.4	19.6	24.7	
	7	Malignant neoplasm of trachea, bronchus and lung	22.5	36.0	27.2	30.0	33.7	38.0	33.7	38.0	20.6	27.0	29.4	32.3	
	8	Malignant neoplasms of breast	33.1	27.6	25.7	25.5	24.1	19.7	25.9	32.1	27.5	26.0	22.1	17.6	
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	16.7	8.7	13.2	11.3	15.5	17.6	11.0	14.8	16.9	20.6	10.3	14.5	
	10	Diseases of the urinary system	7.5	14.5	10.1	16.6	13.1	9.8	11.8	9.6	10.8	11.0	5.1	6.1	
29	Malignant neoplasm of bladder	4.5	..	..	..	..	..	..	..	..	..	..	..		

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Westminster, 2001-2012, <sup>1,2,3,4,5,6,7,8,9,10,11</sup>

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	133.9	141.9	136.4	108.1	98.6	87.8	83.9	81.4	70.4	82.3	58.5	90.4
	2	Cerebrovascular diseases	52.3	68.5	67.0	48.2	46.7	48.7	32.6	27.6	26.8	38.0	30.8	31.3
	3	Malignant neoplasm of trachea, bronchus and lung	59.2	42.8	46.3	60.3	40.4	32.9	42.5	37.1	29.4	48.5	37.0	44.8
	4	Chronic lower respiratory diseases	38.2	28.9	33.8	38.8	38.2	31.7	22.9	27.4	20.3	21.6	27.7	25.9
	5	Influenza and pneumonia	36.9	27.9	36.0	38.4	36.1	20.5	23.5	10.5	23.5	25.2	19.7	13.8
	6	Malignant neoplasm of prostate	19.6	28.6	28.8	22.2	19.2	13.6	22.1	13.7	20.1	18.0	22.7	17.2
	7	Heart failure and complications and ill-defined heart disease	11.5	..	22.8	23.1	29.0	21.6	20.2	18.6	24.8	29.8	14.3	..
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	26.1	12.7	13.7	21.4	15.0	20.6	13.8	10.7	17.3	21.3	13.9	18.4
	9	Cirrhosis and other diseases of liver	25.2	19.8	19.0	22.4	24.3	14.9	15.6	16.0	16.7	12.5	15.8	18.7
	10	Suicide and injury/poisoning of undetermined intent	13.4	12.2	10.4	10	16.9	7.9	12	7.9	6.5	11	11	12
	25*	Malignant neoplasm of bladder	..	..	9.3	..	..	..	..	..	8.8	..	..	..

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	60.8	56.9	66.9	51.8	44.7	35.6	47.3	46.2	37.5	33.3	36.2	26.9
	2	Cerebrovascular diseases	46.2	44.0	52.1	39.9	32.5	40.2	32.1	23.6	23.3	26.9	23.7	22.2
	3	Influenza and pneumonia	23.6	34.7	28.5	27.7	23.7	22.0	18.1	12.3	13.8	21.1	14.1	15.6
	4	Malignant neoplasm of trachea, bronchus and lung	19.8	17.9	35.9	31.2	25.8	19.0	36.3	19.2	25.4	27.1	41.2	18.8
	5	Malignant neoplasms of breast	24.5	30.7	30.6	30.1	26.8	32.7	15.1	20.6	21.3	28.9	16.5	19.0
	6	Chronic lower respiratory diseases	17.8	14.6	23.3	20.3	20.9	21.5	13.5	18.6	11.7	21.5	17.8	17.5
	7	Dementia and Alzheimer's disease	6.2	10.4	12.1	12.3	15.2	..	10.6	8.3	12.2	10.2	15.7	23.3
	8	Heart failure and complications and ill-defined heart disease	16.2	9.8	13.7	8.6	17.1	15.9	17.0	12.7	15.2	16.1	6.7	7.5
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	15.2	12.7	10.9	15.4	14.0	10.4	..	12.5	13.5	7.0	10.5	10.2
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	11.6	9.1	12.2	12.4	10	4.4	18	9	12	7.3	13	10
	30*	Malignant neoplasm of bladder	..	..	..	..	..	..	..	..	..	..	..	..

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: [www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html](http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html)

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 \* Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics