

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Barking and Dagenham, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	244	207	245	176	170	173	131	137	154	146	105	123
	2	Malignant neoplasm of trachea, bronchus and lung	83.4	80.8	73.7	62.7	82.6	73.5	72.2	87.4	74.4	57.9	80.1	59.9
	3	Chronic lower respiratory diseases	55	54.8	68.6	60.8	71.1	49.4	69.6	58.6	63.9	57.9	66.1	55.2
	4	Cerebrovascular diseases	60	60.9	63.1	62.7	54.2	57.2	38.9	53.8	37	32.2	59.7	37.4
	5	Influenza and pneumonia	49.1	43.6	84.3	46.9	53.7	43.1	53.1	41.2	39.8	24.2	32.3	29.4
	6	Malignant neoplasm of prostate	26.9	21.4	22	29.1	29.6	21	30.3	23	21	17	24.7	37.9
	7	Malignant neoplasm of colon, sigmoid, rectum and anus	20	36.2	22	20	27.3	33.2	14	25	18	17	21	15
	8	Aortic aneurysm and dissection	32	23	19	31.1	14	20	25.3	17	20	21	16	..
	9*	Dementia and Alzheimer's disease	12	..	10	13	11	19	..	17	19	18	26.8	34.5
	9*	Malignant neoplasms of lymphoid, haematopoietic and related tissue	21	..	26	22	21	19	17	18	19	23	20	16
18	Malignant neoplasm of bladder	14	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	108	88.9	102	93.3	92	68	62.1	71.6	65.6	52.9	45.6	38.6
	2	Cerebrovascular diseases	64.5	55.7	63.5	42.7	50.5	50.2	52.8	41.5	36.8	35.4	30.5	28.7
	3	Influenza and pneumonia	35.6	50.2	62.9	40.2	39.5	46.1	27	38.8	31.3	28.8	34.7	32
	4	Chronic lower respiratory diseases	33.3	44	48	38.1	54.7	42.7	29.9	37.7	38.7	40.4	30.5	48.7
	5	Malignant neoplasm of trachea, bronchus and lung	44.9	38.7	38.5	33.7	38.8	39	43.3	46.1	50	35.4	36.9	49.1
	6	Dementia and Alzheimer's disease	12.2	17	19.8	17.9	14	12.6	16.3	19.1	21.1	21.7	31	33.4
	7	Malignant neoplasms of breast	31.4	31.7	32.1	30.6	33.2	29.2	32.2	30	33.4	24.1	19.6	25.7
	8	Diseases of the urinary system	16.3	13	13.6	11	17.6	18.7	12.8	24.8	18.9	17.9	8.5	9.5
	9	Heart failure and complications and ill-defined heart disease	9.7	17.9	11.4	14.1	16.9	12.6	10	10	12.1	14	..	8.1
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	14	17	17.4	16	14	17.7	15	11	14	15	13	16
25	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Barnet, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	150	145	131	134	122	123	96.2	111	103	98.9	77.1	72.7
	2	Cerebrovascular diseases	66.2	51.6	52.7	50.5	42.2	37.8	36.6	31.7	40.5	34.8	31.5	34.1
	3	Influenza and pneumonia	42.1	44.4	41.7	36.3	35.3	28.8	30.1	38.8	39.1	34.3	25.9	21.2
	4	Malignant neoplasm of trachea, bronchus and lung	58	42.3	39.9	41.5	36.3	35.7	39.2	30.5	31.9	31	33.1	31.9
	5	Chronic lower respiratory diseases	37.2	30.8	34.8	33	28.3	32	18	26.7	23.5	25.9	22.1	27.3
	6	Heart failure and complications and ill-defined heart disease	24.2	23.6	20.7	18.3	27.6	22.3	19.8	26.1	16.9	26	14.3	19.9
	7	Malignant neoplasm of prostate	22.8	21.7	26.9	19.8	19.1	24.4	25	18.6	21.8	17.7	23.6	21.9
	8	Dementia and Alzheimer's disease	16.3	13.8	12.2	11.6	10.6	12.1	13.9	16	14.1	17.1	26	24.5
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	20	22.5	19.3	22	13	20.5	22.2	18.3	21.2	17.1	19.4	20.3
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	15.9	23.1	20.9	17.2	18.2	18.4	16.7	19.4	14.4	22.5	12	12.9
	19	Malignant neoplasm of bladder	7.6	9.1	8.3	5.4	7.6	5.7	..	6.6	6.8

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	74.7	73	61.3	56.6	50.4	50.5	38.8	46.5	44.9	44.5	39.1	37.3
	2	Cerebrovascular diseases	52.7	46.6	45	37.9	41.9	35.9	32.4	31.9	32.2	31.4	28.8	25.1
	3	Influenza and pneumonia	33.2	32.8	33.6	32.5	35.1	26.5	30.2	32.7	23.7	22.8	22.3	20.1
	4	Heart failure and complications and ill-defined heart disease	22.6	22.3	21.1	24.8	27.4	19.5	17.8	19.7	22.7	23.3	22.5	27.3
	5	Dementia and Alzheimer's disease	12.2	15.1	16.7	17.5	9.7	16.8	12.9	15.7	14.4	20.9	28.3	27.2
	6	Malignant neoplasms of breast	31.1	27.2	33.8	36.5	25.7	18.2	21.1	30.5	23.4	26.9	27.7	21
	7	Malignant neoplasm of trachea, bronchus and lung	17.3	21.3	17.8	23.9	23.7	16.8	20.4	24.2	29.2	24.2	21.8	18.2
	8	Chronic lower respiratory diseases	17	19.6	18	12.4	17.4	15.6	14.8	10.7	12.7	13.1	15.9	16.2
	9	Diseases of the urinary system	10.2	13.3	11.6	11.5	9.8	10.2	9.8	10	11.1	10.1	5.5	7.8
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	14.8	14.2	14.1	21	17.8	11.8	13.9	8.5	12.4	11.3	11.1	12.8
36*	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Bexley, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	170	161	139	152	132	129	118	114	97.6	97.9	100	89.8
	2	Malignant neoplasm of trachea, bronchus and lung	65.5	64.8	50.1	48.9	50	53.3	68.3	45.2	52.1	37.9	56.3	44.6
	3	Cerebrovascular diseases	66.1	71.2	58.4	53	43.4	61.1	49.1	48.1	37	38.8	25.6	26.9
	4	Chronic lower respiratory diseases	49.2	44	44.1	39.4	40.7	34.2	38.3	35.2	35.8	35.1	38.5	25.1
	5	Influenza and pneumonia	30.3	25.4	29.4	23.6	31	25.6	17.8	27.4	18.6	14.7	21.5	29.5
	6	Malignant neoplasm of prostate	38.9	26.3	25.6	34.8	24.3	18.1	22.6	23.2	17.7	19.1	21.7	17.1
	7	Malignant neoplasm of colon, sigmoid, rectum and anus	22.5	15	25.2	22.8	25.5	17.2	17.1	13	25	18.2	21.1	23.2
	8	Dementia and Alzheimer's disease	9.7	16.8	13	13	8	17.4	14.3	16.5	15.7	15.8	18.2	29.7
	9	Heart failure and complications and ill-defined heart disease	15.9	22.2	17.5	13	14	20.9	12	15.2	22.6	16.7	9.6	15.6
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	14	20.7	20.9	21.1	17.4	18.8	9.7	20.5	14.9	15.6	15.7	15.9
18	Malignant neoplasm of bladder	13	12	6.9	7.7	9.6	..	8.4	9.5	11	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	79.9	77.3	78.5	74.5	65.7	56.6	51.1	57.8	46.1	49.7	40.1	41.7
	2	Cerebrovascular diseases	59.4	55.4	72.1	59.7	48.5	44.2	42.1	43.7	29.2	34.7	27.6	20.2
	3	Influenza and pneumonia	34.3	31.1	34.9	23.7	18.9	23.1	23.3	22.7	23.2	19	16.6	15.3
	4	Dementia and Alzheimer's disease	17.6	18.7	20.1	14.2	17.1	15	13.8	13.1	16.2	19.5	31.1	38.6
	5	Malignant neoplasm of trachea, bronchus and lung	22.5	23	33.1	26	28.6	34	33.3	33.5	29.4	35.7	29.2	32.5
	6	Chronic lower respiratory diseases	26	26.6	27.3	23.3	26.8	23.3	18.7	20.9	18.3	18.1	16.4	24.9
	7	Malignant neoplasms of breast	29.8	29.7	34.9	20.4	21	27	25.1	24.1	27	16.8	19.2	17.6
	8	Heart failure and complications and ill-defined heart disease	12.9	11.1	12.5	15.2	13.6	8.1	10.4	18.4	17.2	12	7.4	5
	9	Diseases of the urinary system	6.8	13	14.2	12.8	17.4	15.6	13.7	15	9.3	8.7	8.5	6.2
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	11.7	15.2	10.5	14.9	10.4	11.9	12.2	11.3	14.7	13.3	10.3	10
29	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Brent, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	194	187	156	156	133	123	127	120	107	107	86.1	117
	2	Cerebrovascular diseases	69.1	57.5	57.1	54.2	60.2	48	46.2	44.6	48.1	44.6	33.7	41.8
	3	Malignant neoplasm of trachea, bronchus and lung	48	46.6	43.9	49.5	44.1	50	37.1	47.1	35	45.6	35.2	33.5
	4	Influenza and pneumonia	43.6	41.6	61	46.1	36.5	34.8	39.7	34.4	30.2	35.2	24.7	36.5
	5	Chronic lower respiratory diseases	40.5	53	40.7	33.5	35.6	26.1	28.1	21.8	23.5	20.2	23.8	26.1
	6	Malignant neoplasm of prostate	25	28.4	34.6	35.9	27.1	15	24.2	25.9	32.6	20.7	17.7	18.2
	7	Malignant neoplasms of lymphoid, haematopoietic and related tissue	16.8	22.8	21.5	17.8	16.8	13	20	15	21.4	14.7	16.2	16.3
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	21.7	22.8	23.2	19.4	18.9	12	18.5	12	17.2	23.2	15.2	11
	9	Cirrhosis and other diseases of liver	12	21.6	16	20.9	22.7	29.8	17.2	13	19.3	10	16.6	16.9
	10	Heart failure and complications and ill-defined heart disease	12	18	18.8	21.4	17.7	13	18.6	15	15.1	17.1	9.9	21.7
22	Malignant neoplasm of bladder	7.8	9.2	8.8	9.1	

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	79.1	72	70.5	59.9	63.1	56.6	52.5	54.8	49.8	50.9	50.3	52.3
	2	Cerebrovascular diseases	52.8	39.8	50.5	53.9	43.7	42.9	37.3	38.7	36.6	37.7	33	27.7
	3	Influenza and pneumonia	29.2	28.4	32.9	35.9	36.4	24.8	24.8	29.4	24.1	25.7	15.1	23.2
	4	Heart failure and complications and ill-defined heart disease	13.5	21.2	19.9	22.2	20.1	23.6	12.9	16.2	19.7	24.2	16.2	21.2
	5	Malignant neoplasms of breast	37	32.5	30.5	23.4	28.8	22.8	21	26.5	20.6	22.4	22	20.1
	6	Malignant neoplasm of trachea, bronchus and lung	20.3	17.3	19.2	18.4	15.8	14.9	19.4	21.7	24.9	19.2	22.5	27.3
	7	Chronic lower respiratory diseases	21.5	22.7	17.6	19.7	16.5	11.7	14.1	15	16.4	11.9	7.8	17.8
	8	Dementia and Alzheimer's disease	9.6	9.4	6.5	7.1	8.4	10.5	11	12.5	9.1	13.6	18.4	19
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	15.8	12.8	14.6	18.1	14.4	9.9	9.9	14.4	8.4	12	11	11.2
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	14.3	8	11	11	11	9.5	14.1	10	15.3	14.2	13	7.6
31	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Bromley, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	151	137	127	128	111	102	105	94.7	93.6	90.3	95.1	77
	2	Cerebrovascular diseases	64.9	52.9	46.7	44	40.2	32.9	45.9	37.4	49.1	39.7	28.2	26.5
	3	Malignant neoplasm of trachea, bronchus and lung	42.8	41.6	52	50.9	43.4	47.5	40.6	37.4	38.1	36.9	34.9	33.4
	4	Chronic lower respiratory diseases	38.5	34.7	39.7	36.7	35.5	30.1	29.1	27.7	31.5	24.2	21.9	28.6
	5	Influenza and pneumonia	41.1	37.6	44.2	42.5	31.8	21.8	27	38.5	24.8	19.1	16.5	19.4
	6	Malignant neoplasm of prostate	26.7	23.4	27.7	28.4	31.9	27.2	21.2	26	27.5	26.6	20.1	20
	7	Malignant neoplasm of colon, sigmoid, rectum and anus	29.3	24.8	26.2	25.8	17.6	19.2	22.7	21.2	21.8	21.8	20.1	16.8
	8	Dementia and Alzheimer's disease	10.4	12.3	11.4	6.6	13.1	14.8	18.8	15.1	14	15.3	25.6	30.5
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	24.4	21.7	18.5	19.5	17.3	15	20.8	10.7	13.8	12	12.9	12.8
	10	Heart failure and complications and ill-defined heart disease	9.5	13.1	18.9	16.4	16.2	11.3	11.6	13.9	16.7	14.5	11.4	6.1
18	Malignant neoplasm of bladder	<i>9.1</i>	<i>7.6</i>	<i>5.5</i>	..	<i>9.4</i>	<i>7.7</i>	<i>8</i>	<i>8.6</i>	<i>7.4</i>	..	<i>7.7</i>	<i>10.3</i>	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	68.3	59.1	58.7	55.4	48.3	45.8	47.4	47.5	43.2	33.7	36	34
	2	Cerebrovascular diseases	55.7	40.9	51.5	43.1	47.8	36.3	41.6	31	33.8	33.9	27.8	34.7
	3	Influenza and pneumonia	28.4	40	35.4	29.6	25.7	20.4	25.6	27.6	20.8	14.8	19.7	19.9
	4	Dementia and Alzheimer's disease	18	19.4	14.4	16.5	13.4	15.4	19.2	23.1	23.6	22.1	29.2	28.2
	5	Chronic lower respiratory diseases	19.8	24.7	21	22.9	18.5	19.8	19.9	15.8	12.8	18.3	15.3	22.8
	6	Malignant neoplasm of trachea, bronchus and lung	23.4	20.2	30.8	26.2	23.6	22.8	24.2	20.9	25.8	24.8	23.4	24.4
	7	Malignant neoplasms of breast	28.1	29.3	26.5	27.1	24	27.8	28.1	23.1	20.9	25.8	28	21.1
	8	Heart failure and complications and ill-defined heart disease	15.1	17.8	19.2	16	11.3	10.4	11.1	8.3	12.5	11.7	8.9	6
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	15.6	14.1	13.1	12.8	9.9	14.3	14.2	14.1	9	15.5	11.5	10.9
	10	Diseases of the urinary system	8.2	6.4	10.4	8.8	10.5	10.3	9	9.1	7	13	6.7	9.4
29	Malignant neoplasm of bladder	<i>3.2</i>	..	<i>3.1</i>	..	<i>2.9</i>	<i>2.3</i>	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Croydon, 2001-2012,^{1,2,3,4,5,6,7,8,9,10,11}

		<i>Rate per 100,000 population</i>												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	146	145	148	135	110	122	106	106	101	113	108	99.7
	2	Cerebrovascular diseases	70.2	69.7	62.8	53.4	50.8	48.4	44.9	34.6	32.4	39.1	29.5	25.9
	3	Chronic lower respiratory diseases	48.6	45.2	54.5	50.8	43.7	46.3	39.2	42	35.5	34.3	29.6	41
	4	Malignant neoplasm of trachea, bronchus and lung	40.5	47	45.3	54.5	40.3	44.7	45	41.1	35.7	37.3	41.4	45
	5	Influenza and pneumonia	55	48	45.1	33.3	37.4	27.8	31.4	29.7	35.1	35.9	27.3	26.4
	6	Malignant neoplasm of prostate	24.3	27.6	28.6	29.5	29.6	33	25.8	26.8	13.8	29.8	24.2	20.3
	7	Malignant neoplasm of colon, sigmoid, rectum and anus	24.6	17.9	19	20.2	20.1	18.8	23.7	22.2	20.4	18.4	17.1	21.3
	8	Dementia and Alzheimer's disease	9.2	8.1	9.8	14.2	9	13.5	11.9	12.8	17.9	17.1	28.5	36.5
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	18.4	17.4	12.7	14	18.6	19.3	15.4	17.3	17.2	15	19.8	21.4
	10	Heart failure and complications and ill-defined heart disease	9.6	13.2	14.9	16.1	18.1	17.1	12.7	14.3	23.4	23.1	14.1	5.1
20*	Malignant neoplasm of bladder	9.3	8.9	8.9	9.1	5.9	7	9.8	6.4	8.9	6.1	9.3	..	

		<i>Rate per 100,000 population</i>												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	77.5	85.4	71.5	67.9	61.5	45.9	50.8	48.7	52.4	49.6	46.6	42.3
	2	Cerebrovascular diseases	60.5	51.9	60.5	49.5	48.9	36.7	41	41.1	34.8	36	35.5	27.1
	3	Influenza and pneumonia	49.5	43.9	47.6	40.9	37.7	34.7	23	32.5	31.1	30.2	29.6	25.1
	4	Dementia and Alzheimer's disease	15	13.9	12.4	14.1	10.6	20.2	16.3	22.5	17.2	20.6	31.2	33
	5	Chronic lower respiratory diseases	16	25.2	22.1	26.7	24.6	28	22.6	27.5	24.5	18.5	18.3	29.2
	6	Malignant neoplasms of breast	28.4	36	30.5	36.2	34.7	31.6	23.2	17.2	21.5	23.9	21.2	23
	7	Malignant neoplasm of trachea, bronchus and lung	26.6	28.1	23	27.7	25.2	23.5	27.9	25.2	23.7	19.3	24.3	29.3
	8	Heart failure and complications and ill-defined heart disease	14.1	16.7	12.1	9.1	8.8	15.5	16.7	11.6	10.8	14.8	10.8	7.1
	9	Diseases of the urinary system	10.2	7.2	11.9	12.6	13.3	16.5	10.8	22.5	10.1	10.9	10	9.6
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	11.2	15.5	12.6	12.4	11.7	17	9.8	14.7	11.6	12.2	9.8	10.4
30	Malignant neoplasm of bladder	3.1	5.8	..	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Ealing, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	190	167	189	165	147	134	140	131	120	119	107	90.1
	2	Cerebrovascular diseases	60.8	56.5	61.1	59.8	47.5	54	41.8	35.3	35.7	47.9	30.5	29.4
	3	Influenza and pneumonia	69.7	50.6	56.3	54.3	54.4	41.2	44.1	39	33.4	30.6	34.4	40.2
	4	Malignant neoplasm of trachea, bronchus and lung	52.2	35.5	54.7	50.4	50.1	43.7	50.8	35.3	46.1	33.6	37.5	41.8
	5	Chronic lower respiratory diseases	34.6	43.9	36.3	31.9	37.6	37.2	23.5	28.5	34.7	38.8	27.2	27.3
	6	Malignant neoplasm of prostate	29	31.2	22.5	21.4	33.6	28.1	18.5	21.1	19.1	21.1	23.6	22
	7	Cirrhosis and other diseases of liver	23.2	24.4	26.3	24.3	23.2	17.5	8.8	19.7	24.4	23.1	16.4	18.2
	8	Malignant neoplasms of lymphoid, haematopoietic and related tissue	21.4	15.8	18.7	14	14	22	18.1	15.2	17.8	13.6	17.2	17.1
	9	Heart failure and complications and ill-defined heart disease	12	14.3	17.9	16.6	13.7	13.5	15.2	12	21.1	21.6	17.5	14.8
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	14	18.4	23.2	15.4	13	17	15.6	17	16.2	18.3	11	20
18	Malignant neoplasm of bladder	7.3	8.2	7.5	8.7	11	6.7	8	9	6.3	7.1	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	74.7	80.7	73.6	72.2	63.8	54.6	56.3	52	52	40.9	41.2	36.8
	2	Influenza and pneumonia	49.7	51.7	48.4	48.8	43.6	28.7	32.5	32.3	31.1	25	20.1	20.4
	3	Cerebrovascular diseases	46.5	51.1	44.4	45	36.1	50.1	30.8	40.7	33.4	31.3	25.7	23.8
	4	Malignant neoplasms of breast	26.1	22.8	29.9	25.6	30.2	23	26.2	30.5	32.1	19.3	26.1	26.2
	5	Heart failure and complications and ill-defined heart disease	19.4	15.1	16.4	16.6	17.4	10.4	16.9	21.6	13.9	15.4	19.2	21.8
	6	Dementia and Alzheimer's disease	16.2	14.7	12.1	13.5	11.9	9.7	7.9	16.6	13.6	18.9	24.1	38.9
	7	Malignant neoplasm of trachea, bronchus and lung	28.8	18.2	21	24.4	30.3	26.1	25.9	22.1	30.1	28.1	22.3	17.3
	8	Chronic lower respiratory diseases	20.8	17.8	23.8	24.8	13.4	19.3	18	22.1	17.1	17	24.1	19.1
	9	Diseases of the urinary system	12.3	14.4	16.2	11.6	17.4	10.9	14.3	16.7	17.6	12.3	7.7	8.7
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	11	12.4	7.1	13.8	12.8	13.3	13.3	15.3	18.7	12.2	9.6	9.8
31	Malignant neoplasm of bladder	5	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Enfield, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	168	153	156	148	135	121	116	111	110	97.1	84	93
	2	Cerebrovascular diseases	58.3	72	56.7	53.4	42.4	46.1	45.1	60.8	40	43.9	37.2	28.8
	3	Malignant neoplasm of trachea, bronchus and lung	43.8	45.9	49.4	48.1	41.9	36.4	47.8	50.8	54.8	32.6	48.3	47
	4	Influenza and pneumonia	46.4	65.1	65.8	51.5	42.3	28.4	23.6	24.3	27.1	32.7	35.1	23.2
	5	Chronic lower respiratory diseases	32.1	34.7	32.6	33.1	31.8	29.6	28.7	31.4	34.1	25.5	26	27.6
	6	Malignant neoplasm of prostate	27.2	23.7	24.6	23.4	24.6	22.8	19.1	27.7	27.9	14	22.4	14.5
	7	Heart failure and complications and ill-defined heart disease	26.5	20.1	28.2	21.1	15.3	15.9	25.3	30.8	17.5	24.8	12.3	16.6
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	21.3	22.7	24.4	25.5	24.3	20.5	16.9	20.5	22.8	14.1	19.9	16.4
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	15.1	27.1	24.2	24.1	26.8	17.8	17.2	12	20.9	17	17.8	17.7
	10	Dementia and Alzheimer's disease	15.3	12.9	16.4	13.8	15.9	17.7	12.1	12.6	14.9	18.2	27.1	22.7
18	Malignant neoplasm of bladder	7	..	8	..	6.9	9.9	12	11	6.2	..	6.4	8	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	82.3	74.9	61.3	64.9	67.8	56.6	56.7	55.7	43.7	49.9	34.6	44.5
	2	Cerebrovascular diseases	56.6	57.6	63	51.3	45.8	48.6	37.5	35.6	40	38.6	30.1	31.1
	3	Influenza and pneumonia	40.2	42.5	58.6	38.4	33.5	26.3	18.7	24	27.7	22.6	19.3	23.8
	4	Heart failure and complications and ill-defined heart disease	19.3	26.3	35.4	26.2	24.5	18	21.6	26.7	25.6	21.7	19.8	33.1
	5	Dementia and Alzheimer's disease	18.3	20.2	18.9	19.7	18.1	17	16.3	14.9	22.2	20.9	27.9	30.6
	6	Malignant neoplasms of breast	29.1	24.2	24.8	25.7	31.7	25.7	28.6	34.3	24.1	24.9	23.9	25.7
	7	Malignant neoplasm of trachea, bronchus and lung	27.3	24.6	23.5	23.5	15.4	25.1	20.9	24.9	27.4	26.9	16.8	23.3
	8	Chronic lower respiratory diseases	22.2	18.2	22	14	21.3	16.5	10.6	20.3	15	13.2	19.8	20.5
	9	Diseases of the urinary system	5	12.1	12.2	10.7	12	12.8	12.8	12.5	11.4	14	4.1	4.3
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	16.7	16.2	13.5	15.9	16.6	11.8	11.9	18.4	10.8	9.5	14.5	10.2
	32	Malignant neoplasm of bladder

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Greenwich, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	190	192	212	163	178	162	162	144	148	101	94.5	108
	2	Malignant neoplasm of trachea, bronchus and lung	75.4	69.9	67	72	84.1	74.1	74.3	63.1	72.9	50.7	68.3	52.1
	3	Cerebrovascular diseases	83.7	76.1	78	70.4	70	74.9	54.8	52.4	46.8	45.5	47	44.6
	4	Chronic lower respiratory diseases	69.9	59.8	77.7	60.7	58.8	54.4	63	61	56.4	44.6	50	47.9
	5	Influenza and pneumonia	45.1	47	39.2	32.5	50.4	48.8	33.1	35.5	32.6	31	19.7	16
	6	Dementia and Alzheimer's disease	17	24.3	20.2	21.3	25.7	33.2	26.2	27.5	28.8	21.3	43	44.6
	7	Malignant neoplasm of prostate	21.8	32.6	26.7	21.6	45.2	21.2	29	22	24.9	29.7	24.3	26.1
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	21.5	37.7	27.9	18.3	21.7	15	31.4	18	20.2	18	15	18.5
	9	Cirrhosis and other diseases of liver	21	20.7	21.9	21.6	26.4	33.9	24.9	21	23.3	15	14	16
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	17	20.1	16	22.2	21.4	22.3	15	16	18	13	21	11
	18	Malignant neoplasm of bladder	15	..	11	10	13	11

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	88	74.8	76.9	87.2	82	66	62.6	73.5	55.3	57.3	48.8	44.4
	2	Cerebrovascular diseases	68.1	69.8	80.5	50.7	52.2	46.5	44.3	41	36.8	47.8	33.1	30.9
	3	Dementia and Alzheimer's disease	18.3	18.2	21	25	24.8	19.2	27.7	34.1	26.5	34.6	42.9	61
	4	Chronic lower respiratory diseases	41.5	23.2	38.2	33.2	37.7	25.9	25.9	27.8	31.9	40.9	32.7	39.4
	5	Influenza and pneumonia	30.8	36.6	33.4	24.4	23.9	26.3	21.1	26.7	27.5	26.9	12.5	24.4
	6	Malignant neoplasm of trachea, bronchus and lung	31.4	42.1	39.5	36.6	34.6	35.1	43.7	34.5	26.9	40.3	27.6	31.9
	7	Malignant neoplasms of breast	32.8	34.3	32.7	27	26.8	28.9	23.4	22.3	23.9	25.4	21.9	17.1
	8	Diseases of the urinary system	10.9	18.5	10.5	15	18.8	23.1	11.7	18.2	13	13.3	8.7	12.1
	9	Heart failure and complications and ill-defined heart disease	13.1	13.2	19.3	9.5	7	9.5	9.5	11	8.7	9.4	10.5	..
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	12	11.9	18.4	20.4	19.5	14.2	11	13	14.4	10.9	12	7.9
	28	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Harrow, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	133	158	149	146	124	123	121	104	92.2	97.9	87.7	90.5
	2	Cerebrovascular diseases	71.1	52.5	51.9	42.1	43.4	43.3	34.9	32.8	32.2	30.2	28	26.4
	3	Influenza and pneumonia	40	50.2	33	29.2	30.1	35.6	27.8	32.1	31.4	24.4	36.5	38.5
	4	Malignant neoplasm of trachea, bronchus and lung	52.5	41.4	45.2	48.7	37.6	33.6	23.5	26.8	33.3	30.7	32.6	40.8
	5	Chronic lower respiratory diseases	24.5	32	28.6	35.5	28.6	32.1	25.1	23.9	27.6	17.2	17	20.4
	6	Malignant neoplasm of prostate	24.1	25	16.6	28.2	31.2	30.3	16.2	21	17.7	17.1	17.1	8.6
	7	Heart failure and complications and ill-defined heart disease	13	18	16.1	21.7	17.5	18.3	11	13.6	16.2	13.6	17.8	7.9
	8	Malignant neoplasms of lymphoid, haematopoietic and related tissue	25.4	25.3	13	12	21.8	19.2	20.6	15.3	11	13	15	13.6
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	17	18	25.6	19.3	11	17.8	20.6	18.3	11	11	14	16.6
	10	Dementia and Alzheimer's disease	16.6	9.9	12	7.9	9.9	9.8	..	6.4	12	7.4	13.3	11.9
	17*	Malignant neoplasm of bladder	..	9.4	9.1	11	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	68.8	61.7	65.7	64.7	48.5	45.9	58.4	49.6	45.9	37	43.3	38.2
	2	Cerebrovascular diseases	41.1	40	48.2	41.3	42.3	34.8	30.3	31.2	32.4	22.4	19.9	29.8
	3	Influenza and pneumonia	31.1	27.3	30.4	33.5	33.2	25.4	28.9	24.9	29.3	25.1	18.7	23.5
	4	Heart failure and complications and ill-defined heart disease	8.8	22.9	23.4	25.5	24.2	22.9	17.3	21.8	26.7	21.7	20.2	22
	5	Malignant neoplasms of breast	28.9	29	31.4	28	31.2	27.8	28.9	27.9	16.7	30.9	26.5	25.3
	6	Dementia and Alzheimer's disease	16	12.4	12.3	11.3	8	11.7	12.1	9.4	10.2	10.2	14.6	17.4
	7	Malignant neoplasm of trachea, bronchus and lung	16.3	16.1	20.8	13.1	26.1	16.5	20.3	17.2	12.3	19.2	16.5	16.7
	8	Chronic lower respiratory diseases	17.4	18.3	16	10	11.9	11.1	11	12.3	16.1	12.6	11.5	14
	9	Diseases of the urinary system	10.3	7.2	9.2	10	11	11.2	7.7	8.2	5.8	8.3	10.5	5.5
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	11	14.2	12.2	11.9	13	11	13.8	12	8.5	10	5.8	8.5
	32*	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Havering, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	184	192	170	158	142	120	141	127	111	114	88.2	92.1
	2	Malignant neoplasm of trachea, bronchus and lung	69.5	59.5	57.3	56.3	55.6	56.3	38.1	48.9	44.5	55.7	40.8	43.1
	3	Cerebrovascular diseases	67.8	64	56.4	54.3	44.8	59.3	41.3	45	34.2	45.2	34.2	30.3
	4	Chronic lower respiratory diseases	41.7	49.6	59.5	44.7	44	36.6	43.8	46.1	33.7	30.2	38.8	38.2
	5	Influenza and pneumonia	53.4	45.5	36.2	43.8	32.5	37.2	31.6	40.3	44.4	28.7	19.2	32.8
	6	Malignant neoplasm of prostate	31.2	32.4	25.6	27.6	22.2	24.8	27.4	24.1	22.5	26.8	24.7	24.3
	7	Malignant neoplasm of colon, sigmoid, rectum and anus	20.8	22.9	25.6	26.9	23.7	15	22.6	17.8	31.6	16.7	18	13.7
	8	Dementia and Alzheimer's disease	17.6	17.1	7.3	11	11	12	16.9	8.6	23.2	17.3	31.6	36.7
	9	Aortic aneurysm and dissection	22.1	22.9	21.7	18.4	15.6	21.2	11.6	11	14.6	14.4	10	15.5
	10	Heart failure and complications and ill-defined heart disease	10	17.5	21.5	17.3	18.7	17.9	12.3	12.1	16.1	13.9	..	8.9
17	Malignant neoplasm of bladder	..	11	8.3	11	11	7.5	6.9	9.1	8.8	9.1	6.9	..	

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	85.5	94.3	82.1	66.7	71	51.3	56	52.7	45.3	46.7	32.6	44.7
	2	Cerebrovascular diseases	53	56.9	57.6	45.2	52.1	38.7	30	37.3	41.2	31.1	29.9	27
	3	Influenza and pneumonia	38.5	39.6	41.5	37.3	35.5	39.7	30.9	36.4	26.8	28.1	24.3	19.9
	4	Dementia and Alzheimer's disease	21.4	20.9	21.5	17.5	13.4	11.9	16.6	20.2	19.9	25	34.2	45.7
	5	Chronic lower respiratory diseases	20	26.7	32.5	28.7	28.1	29.9	20.2	23.3	20.6	18.6	18.4	32.2
	6	Malignant neoplasm of trachea, bronchus and lung	26.8	25.6	27.7	25.8	33.6	25.6	36.8	32.4	30	29.3	30.2	31
	7	Malignant neoplasms of breast	42.4	34.7	29.7	31	30.4	26.2	26.7	24.2	31.3	13.7	23.8	24.5
	8	Heart failure and complications and ill-defined heart disease	10.8	11.2	17.1	11.5	12.7	15.2	11.1	8.5	7.4	7.8	5	6.1
	9	Diseases of the urinary system	9.5	9.4	11.6	15	12.6	14.8	11.6	11.4	13.7	11.4	6.1	6.2
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	15.1	16.1	14.8	11.2	18.5	11.7	13.5	14.9	13.1	13	11.5	10.4
25*	Malignant neoplasm of bladder	..	4.5	5.7	4.5	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for

2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Hillingdon, 2001-2012,^{1,2,3,4,5,6,7,8,9,10,11}

		Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	179	181	158	142	148	137	142	113	90.8	91.5	101	85.9
	2	Malignant neoplasm of trachea, bronchus and lung	71.6	60.2	50.2	59.8	37.3	59.1	48.8	52.7	35.7	49.5	41.2	43.1
	3	Influenza and pneumonia	47.2	50.9	57.4	61.2	45.4	47.3	36.2	40	32.6	30.7	39.2	26.6
	4	Cerebrovascular diseases	56.5	50.7	53	59.1	44.7	48.1	34.5	36	31.1	31.3	28.7	34.4
	5	Chronic lower respiratory diseases	38.4	38.7	56.8	31.7	33.5	26.5	21.7	41.2	33.4	30.9	30.2	36
	6	Malignant neoplasm of prostate	24.2	29.7	19.7	23.8	21.6	23.3	20.1	29.8	24.5	25.8	25.9	32.8
	7	Malignant neoplasm of colon, sigmoid, rectum and anus	26.2	25.1	24.4	12	19.3	22.3	22.3	21.3	25.4	16.9	9.8	19.6
	8	Dementia and Alzheimer's disease	19.2	16	13	..	14	8.7	11	14.5	17.1	17.9	24.3	25.7
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	17.2	16.1	17.6	16	17	20.7	13	18.4	22.8	15.3	17.6	15.7
	10	Cirrhosis and other diseases of liver	17.7	17	23.3	26.2	15	29.1	17.8	17.2	16.5	16.9	10	22.9
	17	Malignant neoplasm of bladder	8.6	11	..	12	12	8.1	7.6	..	6.5	9.3

		Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	80	76.8	74.6	62.4	51.1	55.6	63.5	57.2	38.1	37.4	40.4	40.1
	2	Cerebrovascular diseases	55.3	47.3	58.7	46.6	38.4	32.9	30.5	29.9	31.3	37	26.3	27.4
	3	Influenza and pneumonia	37.5	35.7	39.7	46.8	40.5	33.8	27.6	31.3	20.6	23.5	23.1	25.3
	4	Chronic lower respiratory diseases	22.1	20.3	29.2	32.7	27.5	17	31.3	27	22.6	26.6	26.4	26
	5	Malignant neoplasm of trachea, bronchus and lung	27.1	34	31.3	25.2	34.6	28.7	30.7	27.7	31.7	29.5	21.4	40.1
	6	Heart failure and complications and ill-defined heart disease	11	11.6	12.5	13.3	17.6	17.2	19.4	15.8	22.9	24.2	20.8	22.1
	7	Malignant neoplasms of breast	34.1	33.7	30.2	37.3	35.5	19.5	22.9	23.9	30.5	25	22.1	30.2
	8	Dementia and Alzheimer's disease	14.5	21	15.9	16	8.8	7.8	11.5	14	16.4	12.9	24.9	29.7
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	8.6	16.3	8.9	17.2	12.7	12.5	15.4	12.2	12.4	10	11.9	15.3
	10	Diseases of the urinary system	6.8	5.2	12.5	6.6	10	14.7	9	8.6	11.9	9.7	5.6	9.6
	27*	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Hounslow, 2001-2012,^{1,2,3,4,5,6,7,8,9,10,11}

		Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	192	184	150	161	146	121	136	117	105	114	96.8	85.2
	2	Cerebrovascular diseases	65.2	73.5	73.8	59.9	57.1	47.8	50.8	43.4	47.7	37.4	31.2	32.8
	3	Malignant neoplasm of trachea, bronchus and lung	44.9	65.6	44.9	53.6	48.2	60	51.9	48.1	57.9	43.4	38.8	44
	4	Influenza and pneumonia	64.2	66	49.8	65.7	55.6	38.3	35.5	36.6	30.3	32.1	32.2	38
	5	Chronic lower respiratory diseases	46.6	50.9	50	42.3	39.2	47.6	42.3	38.2	25.5	33.6	38.8	43.6
	6	Malignant neoplasm of prostate	24.6	16	23	31.6	27.3	21.9	18.9	29.2	23.5	33.1	22.5	20.9
	7	Cirrhosis and other diseases of liver	27	29.3	36.8	20.7	15	19.6	21.7	26.5	17	19	18	18.2
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	31.8	31.1	26.6	9.8	16	15	19.5	27.1	19.4	17	20	21.8
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	18	21.6	24.7	18	14	15	25.9	12	17	14	15	20.7
	10	Heart failure and complications and ill-defined heart disease	22.3	9.1	28.8	20	31.9	12	19	15	16	20.9	..	13
	18*	Malignant neoplasm of bladder	9.3	

		Rate per 100,000 population												
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	93.9	79.5	90.9	79.3	69.5	64.1	47.9	52.7	48.2	48.5	44.7	38.9
	2	Cerebrovascular diseases	60.6	61.4	44.8	60.6	51	47	40.1	42.5	32.6	32	36.8	29.9
	3	Influenza and pneumonia	51.7	45.4	62.1	56.8	35	28.7	34.1	30.4	26.2	23.4	20.9	28.3
	4*	Chronic lower respiratory diseases	30	32.1	26.8	23.5	25.5	25.5	23.9	16.8	19.8	25.4	25.8	19.7
	4*	Malignant neoplasms of breast	38.8	36.2	36.5	33	31.1	26.5	20.3	19	36.5	26.2	33.6	29.9
	6	Dementia and Alzheimer's disease	20	13.7	19.4	13	16.2	9.9	13	21.8	18.2	16.8	31.3	32.4
	7	Malignant neoplasm of trachea, bronchus and lung	26.8	29.3	26.9	16.5	27	35.3	37.8	28.7	24.1	27.8	29.7	28.8
	8	Heart failure and complications and ill-defined heart disease	23.4	22.5	27.2	17.4	20.1	17.6	16.8	19.2	17.3	11	11.2	9
	9	Diseases of the urinary system	6.6	11.9	13.1	18.5	11	12.1	14.1	18.1	9.5	14.3	8	9.3
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	11	12	12	13.9	8.5	13	13	13	13.4	13	6.5	7.7
	33	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Kingston upon Thames, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	146	146	123	133	117	118	103	83.8	98.6	80.2	91.3	85.4
	2	Cerebrovascular diseases	71.1	58.8	74	59.1	50.2	45.5	44.3	38.1	32.5	38.9	31.9	18
	3	Influenza and pneumonia	49.7	50.5	47.8	37.7	38.9	34.3	37.2	45.1	23.4	31.7	23.2	34
	4	Malignant neoplasm of trachea, bronchus and lung	66.2	41.9	51.5	52.6	34	56.4	42.1	42.2	39.4	35.7	40.3	31.2
	5	Chronic lower respiratory diseases	40.5	28.6	34.5	23.9	27	27.1	36.8	31.3	24.6	25.6	27.3	36.5
	6	Malignant neoplasm of prostate	17	26.6	30.5	33.2	33.4	19	21	20	28.2	22.1	23.1	15
	7	Dementia and Alzheimer's disease	16	12	13	13	12	..	13	16	11	17	26.5	26.1
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	22	20	21	18	23	26.6	..	15	14	17	15	20
	9	Heart failure and complications and ill-defined heart disease	17	25.4	19	..	22.3	19	..	17	..	15	14	..
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	19	21	13	..	21	17	15	13	20	15	16	..
17*	Malignant neoplasm of bladder	13	11	15	13	..	

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Cerebrovascular diseases	67.1	54.6	61.4	58.6	44.5	38.9	49	34.5	44.5	38.7	35.3	30.9
	2	Ischaemic heart diseases	51.5	64.1	66.8	53.6	47.1	46	49.9	50.3	46.7	38.5	35.5	30.1
	3	Influenza and pneumonia	50.5	48.7	28.1	41.4	35.9	41.2	27.7	35.6	25.2	26.5	25.3	22.6
	4	Dementia and Alzheimer's disease	14.9	14.5	18.1	13.9	8.8	15.4	13.4	12.4	12	17.6	28.2	30.8
	5	Chronic lower respiratory diseases	23.6	12.8	18.3	17	19.6	16.5	30.2	24.8	11	15.7	19.3	23
	6	Malignant neoplasms of breast	20	39.1	31.1	24.1	18.4	24.3	22.3	22	19.1	27.3	18	22.8
	7	Malignant neoplasm of trachea, bronchus and lung	24.8	21	18	22.1	30.8	18.9	30.7	23.9	21.9	20.4	24.8	23.5
	8	Heart failure and complications and ill-defined heart disease	16.9	27	16	19	13	12.5	15.3	14.8	7.9	7.3	..	7.8
	9	Diseases of the urinary system	5.1	9.7	8.7	10	8.9	7.7	8.3	11.1	14.2	8	9.7	7.7
	10	Malignant neoplasm of colon, sigmoid, rectum and anus	22.2	17	15	17	16	15	12	9.5	15.8	9.9	9.2	13
28	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

[4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Merton, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	162	128	147	129	84.9	88.6	101	113	91.2	91.9	90.3	82.5
	2	Cerebrovascular diseases	68	63.2	71.2	44.1	54.9	42.2	31.7	43.8	37.5	43.3	33.3	31.9
	3	Malignant neoplasm of trachea, bronchus and lung	49.4	52	43	36.2	50.6	33.5	41.8	38.2	25.9	38.5	33.1	44.6
	4	Influenza and pneumonia	55.1	43.9	39.4	35.3	32.8	39.5	29.8	30.8	30.9	27.8	18	29.4
	5	Chronic lower respiratory diseases	48.7	44.6	58.1	25.9	29.8	32.6	28.7	26	32.2	26.6	26.1	25.8
	6	Heart failure and complications and ill-defined heart disease	20.9	23	32.9	31.2	38.7	33.5	29.4	24.7	29.6	24.7	20.5	13
	7	Malignant neoplasm of prostate	27	21.8	17	28.4	29.5	31.1	13	22.6	23.2	27.7	20.7	27.8
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	24.3	24.2	25.5	20	14	31.4	21.9	20	23.2	21	19	19.6
	9	Malignant neoplasms of lymphoid, haematopoietic and related tissue	15	19	23	24	17	19	23.8	15	18	14	22.8	12
	10	Dementia and Alzheimer's disease	15	..	15	..	12	11	14	19	9.1	14	32.9	23.8
17*	Malignant neoplasm of bladder	13	..	13	

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	65.3	66.8	66.1	59.7	53.6	52	47.1	50.8	54.5	37.8	38	41.2
	2	Cerebrovascular diseases	53.2	48	56.2	47.5	46	46.5	33.6	44.1	27.9	32.7	31.2	30.3
	3	Influenza and pneumonia	32.3	34.7	33.8	24.6	25.8	21.1	27.2	29.2	21.4	16.8	22.4	19.8
	4	Malignant neoplasms of breast	35.8	30.5	30.7	32.5	29.8	34.7	27.4	22.8	23.2	26.8	20.7	24.7
	5	Chronic lower respiratory diseases	22.7	24.2	25.7	23.8	25.2	21.2	18	20.7	23.5	15	24.6	23.8
	6	Dementia and Alzheimer's disease	18	10.7	13.4	10	10.5	5.2	9.8	14.2	15.1	20.1	27.2	31.7
	7	Malignant neoplasm of trachea, bronchus and lung	35.9	21.7	26.4	26	26.9	22.1	27.6	23	29.4	18.1	24.9	32
	8	Heart failure and complications and ill-defined heart disease	14.1	11.2	22	21.1	15.1	16.8	15.4	12.2	16.6	16.3	10	8.8
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	11	16	14.2	15	20.2	15.7	17.3	18.1	14	6.8	20.4	10
	10	Diseases of the urinary system	8.5	10	10.6	10	11	11	6	12.2	11	10.4	..	9.6
27	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Redbridge, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	207	164	181	137	166	116	117	123	116	111	98.3	92.9
	2	Cerebrovascular diseases	52.7	57.6	50.4	63.4	57.6	41.1	41.1	52.1	47.1	43.7	35.5	32.7
	3	Influenza and pneumonia	53.5	49.2	59.7	45.6	42.4	33.7	34.9	21.8	32.3	32.7	24.2	36.6
	4	Malignant neoplasm of trachea, bronchus and lung	58.1	55.2	37	47.1	26.2	46.5	43.6	45.1	30.5	44.6	30.2	30.6
	5	Chronic lower respiratory diseases	35.3	31.4	28	42.5	35.3	27.4	32.3	26.3	26.7	27.9	26.3	24.5
	6	Malignant neoplasm of prostate	27.1	27.7	22.8	17.8	35.3	20.5	21.8	18.1	20	22.4	22.1	17.6
	7	Dementia and Alzheimer's disease	15.2	20.4	15.1	7.6	15.4	16.2	13.2	14	14.2	17	27.5	26.2
	8	Malignant neoplasms of lymphoid, haematopoietic and related tissue	20.3	22.3	22.4	21.7	14	18.9	17.6	19.4	13	15.1	23.3	11
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	21.6	19.5	23.4	15	15	12	19.1	15	15.9	14	15.7	17.4
	10	Diseases of the urinary system	15.6	15	13	14.9	11	16.1	10	13.7	15.4	14.7	13	10
	17	Malignant neoplasm of bladder	8.3	9.1	7.8	11	8.6	7.4	9.8	8.5	8.6	6.2

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	78.2	87.2	89.5	68.6	73.8	69.5	62.4	64.5	54	48.5	38.6	39.3
	2	Cerebrovascular diseases	56.8	48.6	56.2	63.4	50.4	27.9	29.5	48.3	35.8	35.5	32.9	32.8
	3	Influenza and pneumonia	43	35.5	39.9	35.9	35.9	32.6	28.3	27.2	22	23.5	24.1	33.9
	4	Dementia and Alzheimer's disease	13.9	16.7	19.4	16.8	14.3	18	15.5	21.9	26	25.7	27	38.9
	5	Malignant neoplasms of breast	31.8	36.2	30	23.7	24.9	28.3	28.7	28.3	26.6	25.7	25.5	22.7
	6	Malignant neoplasm of trachea, bronchus and lung	26.2	30.6	26.9	19.4	25.3	20.6	17.5	19.8	30.8	22.9	20.7	17
	7	Chronic lower respiratory diseases	18.3	18.4	17.9	19.1	14.6	15.6	18.8	16.8	16.2	16.6	18.1	13.5
	8	Diseases of the urinary system	12.5	8.7	16.3	15.1	14.2	12.3	10.8	13.7	10.4	11.4	10.3	8.5
	9	Heart failure and complications and ill-defined heart disease	8.2	8.5	12.3	9.8	7.9	10.4	13.6	8.5	7.9	10.7	5.2	8.1
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	12.2	17.7	11	9.8	17.5	12.9	11.6	9.2	12.5	13.9	15.6	13
	27*	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Richmond upon Thames, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	111	133	116	128	117	97.5	96.5	89.2	84.7	77.6	60.6	67.4
	2	Cerebrovascular diseases	67.8	56.5	58.4	53.1	36.9	41	41	35.2	50.2	32.9	22.4	27.7
	3	Influenza and pneumonia	54.7	51	58	41.5	34.6	35.2	33.1	35.9	27.8	22.6	33.2	28.8
	4	Malignant neoplasm of trachea, bronchus and lung	39.9	39.3	58.7	35.9	34.4	42.2	39.1	34.3	26.4	40.8	35.3	41.8
	5	Chronic lower respiratory diseases	32.6	23.9	22.9	34.2	26.8	26.6	25.2	34.7	21.1	33	32	33.3
	6	Malignant neoplasm of prostate	27.9	25.9	16	34.8	19.4	24	26.3	24.3	16	24.1	25.9	17
	7	Malignant neoplasms of colon, sigmoid, rectum and anus	18	21.7	15	16	16	20	21.8	17	..	20	18.9	25.4
	8	Malignant neoplasms of lymphoid, haematopoietic and related tissue	16	19	23.3	24.9	14	16	17	11	15	17	20.1	20.9
	9	Heart failure and complications and ill-defined heart disease	18	17	19.6	17	11	21.4	13	17	12	16.9
	10	Dementia and Alzheimer's disease	12	11	9.1	12	11	11	..	35.2	26.2
15	Malignant neoplasm of bladder	10	14	11	..	9	9.2	

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	59	51.3	63.7	47.9	45.7	52.4	35.9	42.4	34.1	25.9	31.1	33.3
	2	Cerebrovascular diseases	42.7	43.7	48.9	40.3	40.1	38	37.1	34.7	25.3	32.4	22.3	21.1
	3	Influenza and pneumonia	37.6	45.2	37.9	29.7	26.5	20.9	23.4	21.2	22.8	16.6	16	18.1
	4	Malignant neoplasm of trachea, bronchus and lung	28.3	24	36.5	32.3	23.9	32	28.9	29.1	21.2	23	19.2	28.5
	5	Dementia and Alzheimer's disease	8.3	11.2	10.5	10.6	14.4	11.2	13.6	9.8	11.8	13.2	24	30.4
	6	Malignant neoplasms of breast	39.1	26.2	23.2	32.5	28.2	37.5	19.8	24.3	22	30.2	23.2	15.7
	7	Chronic lower respiratory diseases	17.2	21.9	24.4	19.2	17.8	18.9	16.7	15.8	9	21.6	17.4	15
	8	Heart failure and complications and ill-defined heart disease	10.7	13.1	13.5	14.6	17.1	12.7	10.2	8.3	8.3	5.4	8.7	5.7
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	15.3	14	12	15	13.7	16.5	14	10	14.7	13	18.5	7.9
	10	Diseases of the urinary system	8.7	10.2	14.2	11	11.8	7.1	7.8	6.3	6.5	11.5	4.4	6.2
34	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Sutton, 2001-2012, ^{1,2,3,4,5,6,7,8,9,10,11}

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	156	149	146	126	110	95.9	99.1	79.3	84.9	82.1	78	62.3
	2	Cerebrovascular diseases	55	55.2	57.7	48.2	63.9	54	43	45.1	36.1	30.4	36.4	39.2
	3	Malignant neoplasm of trachea, bronchus and lung	65.7	43.6	53.7	41.4	49.9	45.5	50.9	42.6	58.4	40.2	36.8	50.3
	4	Influenza and pneumonia	59.9	41.7	34.7	37.6	34.2	39.9	37.2	39.8	35.5	28.1	31.3	25.2
	5	Chronic lower respiratory diseases	51.2	38.5	47.3	26.8	22.3	26.9	33.9	30.2	37.4	31.6	40.7	34.7
	6	Heart failure and complications and ill-defined heart disease	22.4	38.2	36.3	34.1	46.7	32.2	33.1	47.6	32.3	33.2	20.7	7.8
	7	Malignant neoplasm of prostate	23.2	20.8	30.8	21.7	21	19.1	23.9	25.2	22.6	32.4	18	19.5
	8	Malignant neoplasm of colon, sigmoid, rectum and anus	21.9	18	26.6	20.5	20	18	19.6	19.1	17	18	17.1	16
	9	Dementia and Alzheimer's disease	..	10	14	10	11	12	15	14	16.1	23	18.9	30.6
	10	Malignant neoplasms of lymphoid, haematopoietic and related tissue	21.8	13	18	22.8	18	15	23.1	14	16	14	13	20.5
18	Malignant neoplasm of bladder	11	12	9.2	9.4	8.3	..	

			<i>Rate per 100,000 population</i>											
Sex	Rank	Group Description	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	81.3	74.3	79	85	62.3	51.9	57.8	53.4	52.9	39.2	33.1	31.5
	2	Cerebrovascular diseases	66.4	60.1	80.7	55.8	46.5	47.8	40.8	42.6	33.9	30.9	19.4	34.5
	3	Influenza and pneumonia	47.4	38.8	32.9	40.1	32.4	33.4	29.1	32.1	22.5	19.5	21.9	19.8
	4	Dementia and Alzheimer's disease	12.5	11.4	15.1	16.6	15.3	21.3	16	21.2	16.5	18.9	32.1	34.4
	5	Chronic lower respiratory diseases	27.4	31.4	23.3	17.7	14.3	23.9	25.1	16.1	22.2	20.9	24.2	25.3
	6	Malignant neoplasm of trachea, bronchus and lung	33.2	27.8	34.9	20.6	20.6	22.1	26.9	35.7	23.5	40.3	22.5	21.9
	7	Malignant neoplasms of breast	32.3	31.9	35.7	23.2	28.3	31.6	27	24.1	23.8	18.6	22.1	31.1
	8	Heart failure and complications and ill-defined heart disease	13.6	22.2	15.9	21.8	19	15.7	16.6	16.4	13.2	13.8	11	7.7
	9	Malignant neoplasm of colon, sigmoid, rectum and anus	15.7	18.7	14.9	19.4	6.5	15.7	16.3	11	11	13	14	11
	10	Diseases of the urinary system	5.4	6.5	9.1	9.5	7.7	12.4	9.2	14.4	10	12.4	6.8	5.9
26	Malignant neoplasm of bladder	

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 [Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision \(ICD-10\) codes in Health Statistics Quarterly 28, Winter 2005, Annex A](#)

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9 are coded to U50.9).

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics

Age Standardised Rates for the Top 10 leading causes of death including Malignant neoplasm of bladder, by sex, Waltham Forest, 2001-2012,^{1,2,3,4,5,6,7,8,9,10,11}

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Males	1	Ischaemic heart diseases	231	230	204	188	148	159	145	123	134	101	87.8	110
	2	Cerebrovascular diseases	67.5	72.4	61.6	72.9	50.6	59.1	46.3	58.2	46	47.7	34	30.1
	3	Malignant neoplasm of trachea, br	65	42.1	53.4	52.3	53.2	44.2	57.3	47.2	49.1	52	46.6	44.1
	4	Influenza and pneumonia	72.4	64.1	72.5	52.1	70.1	49.9	34.2	36.7	30.1	30.2	25.6	28.9
	5	Chronic lower respiratory diseases	39.9	36.7	37.2	37.5	57	41.7	51.5	32.8	50.2	39	31.7	44.4
	6	Malignant neoplasm of prostate	32.2	28.1	22	27.3	22.2	22	21.6	9.4	19.4	29.3	32.4	27.2
	7	Malignant neoplasm of colon, sigm	16	23	20.8	20.1	20	36.9	13	19	19	13	19	19.8
	8	Dementia and Alzheimer's disease	16	10	20	20	13	20.1	12	18	9.3	17	30.2	31.9
	9	Malignant neoplasms of lymphoid,	18	22.6	16	18	20.6	20.8	21.3	13	17	14	15	12
	10	Cirrhosis and other diseases of liver	11	17	13	17	18	13	21.8	23.5	21.5	16	13	16
	19	Malignant neoplasm of bladder	9.9	18	10	11	11	9.5

Sex	Rank	Group Description	Rate per 100,000 population											
			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Females	1	Ischaemic heart diseases	113	105	96.2	72.4	80.9	66.7	65.4	71.1	67.9	47.5	50.7	42.4
	2	Influenza and pneumonia	46.8	43.5	70.2	49	43	34.6	41.5	32.1	30.5	24.9	28.6	24.6
	3	Cerebrovascular diseases	62.9	52.7	57.2	45.2	46.7	42.2	44	33.5	42.4	33.9	33.1	29.7
	4	Dementia and Alzheimer's disease	13	18.1	19.9	20.7	20.2	16	17	29.5	17.2	19	34.4	31.1
	5*	Malignant neoplasm of trachea, br	39.9	32.5	28.2	26.5	24.1	28.9	35.8	16.5	26.5	27.2	22	31.2
	5*	Malignant neoplasms of breast	33.8	34.4	31.5	28.2	27.3	29.6	16	22.7	29.8	33.8	22	24.9
	7	Chronic lower respiratory diseases	22.8	18.4	13.6	31.2	22.3	25.7	24.2	21.1	18.4	19	24.4	30.4
	8	Heart failure and complications anc	13.8	18	14.7	11.1	10.2	9.1	14.4	17.8	9.8	13.8	7.4	7.7
	9	Malignant neoplasm of colon, sigm	13	21	17.5	21.9	13	17.3	8.6	15.7	10	..	14	8.3
	10	Diseases of the urinary system	6.6	8.9	11.8	15.1	9.8	10	14.4	13.1	15.5	7.2	6.9	10
	27	Malignant neoplasm of bladder	7.2

1 Based on boundaries as of August 2013

2 Figures are for deaths registered, rather than deaths occurring in each calendar year. Due to the length of time it takes to complete a coroner's inquest it can take months or even years for deaths from certain causes to be registered. More information on registration delays for a range of causes can be found on the ONS website: www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/impact-of-registration-delays-on-mortality-statistics/index.html

3 Figures exclude deaths of non-residents

4 Underlying cause of death was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes in Health Statistics Quarterly 28, Winter 2005, Annex A

5 Suicide was defined using the International Classification of Diseases, Tenth Revision (ICD-10) codes X60-X84 and Y10-Y34 for 2001 onwards (excluding Y33.9 where the coroner's verdict was pending for the years 2001 to 2006. From 2007, deaths which were previously coded to Y33.9

6 Suicide figures and rates are for persons aged 15 years and over. Suicide rates are expressed per 100,000 population aged 15 years and over.

7 Rankings are based on the grand total number of deaths for that cause of death

8 Age-standardised rates per 100,000 population, standardised to the 1976 European Standard Population. Age-standardised rates are used to allow comparison between populations which may contain different proportions of people of different ages.

9 .. Rates were not calculated where there were less than 10 deaths in a year

10 Rates based on less than 20 deaths have been italicised and must be interpreted with caution

11 * Where multiple causes of death had the same total number of deaths they share the same rank

Source: Office for National Statistics